

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 July 2001 (05.07.2001)

PCT

(10) International Publication Number
WO 01/47955 A2

(51) International Patent Classification⁷: **C07K 14/00**

(21) International Application Number: **PCT/GB00/04984**

(22) International Filing Date:
22 December 2000 (22.12.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9930294.5 23 December 1999 (23.12.1999) GB
0025234.6 14 October 2000 (14.10.2000) GB

(71) Applicants (for all designated States except US): **MEDICAL RESEARCH COUNCIL** [GB/GB]; 20 Park Crescent, London W1N 4AL (GB). **INTERNATIONAL AIDS VACCINE INITIATIVE** [US/US]; 27th floor, 110 William Street, New York, NY 10038-3901 (US). **UNIVERSITY OF NAIROBI** [KE/KE]; P.O. Box 47970, Nairobi (KE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HANKE, Tomas**

[CZ/GB]; MRC Human Immunology Unit, Institute of Molecular Medicine, The John Radcliffe, Oxford OX3 9DS (GB). **MCMICHAEL, Andrew, James** [GB/GB]; MRC Human Immunology Unit, Institute of Molecular Medicine, The John Radcliffe, Oxford OX3 9DS (GB).


(74) Agent: **KEITH W NASH & CO.**; 90-92 Regent Street, Cambridge CB2 1DP (GB).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: **IMPROVEMENTS IN OR RELATING TO IMMUNE RESPONSES TO HIV**



(57) Abstract: Disclosed is an immunogen in sterile form suitable for administration to a human subject, the immunogen comprising: at least a portion of the gag protein of HIV, said gag protein being from an HIV clade or having a consensus sequence for one or more HIV clades, and comprising at least parts of p17 and p24; and a synthetic polypeptide comprising a plurality of amino acid sequences, each sequence comprising a human CTL epitope of an HIV protein, and wherein a plurality of HIV proteins are represented in the synthetic polypeptide, said CTL epitopes being selected to stimulate an immune response to one or more HIV clades of interest.

WO 01/47955 A2



Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Title: Improvements in or Relating to Immune Responses to HIV**Field of the Invention**

This invention relates to an immunogen designed to elicit an anti-HIV immune response in a human subject (especially a cell-mediated response), a nucleic acid molecule encoding the immunogen, compositions comprising the immunogen and/or the nucleic acid molecule, and to a method of inducing an anti-HIV immune response (especially a cell-mediated response) in a human subject.

Background of the Invention

Development of effective human immunodeficiency virus (HIV) vaccines is one of the primary goals of current acquired immunodeficiency syndrome (AIDS) research. Despite progress in prevention and powerful drug combinations to treat HIV infection, an estimated 16,000 people become infected every day. Over 90% of new infections occur in developing countries for which the recent medical advances are not immediately applicable or affordable. The best hope for these countries is the development of an effective, accessible HIV vaccine. There is now growing optimism among scientists that an AIDS vaccine may be possible (McMichael & Hanke 1999 Nat. Med. 5, 612-614; Gold 1999 IAVI Report 4, pp 1-2, 8-9, 15-16 & 18).

An ideal prophylactic vaccine should induce sterilizing immunity, so that after exposure, the virus would never be detected in the body. However, this is probably an unrealistic objective. Rather, an attainable goal may be a vaccine-induced immunity that results in a limited and transient virus replication, after which the virus becomes undetectable, there are no signs of disease and no transmission to other individuals. Alternatively, a potentially successful vaccine may induce immune responses that at least hold the virus in check at levels so low, that both progression to AIDS and transmission are entirely or substantially prevented.

To induce sterilizing immunity, a prophylactic vaccine may need to elicit both humoral and cell-mediated immune responses. Since HIV was isolated and sequenced, there has been a considerable effort to develop envelope-based vaccines inducing neutralizing antibodies (nAb). However, this has proved to be exceedingly difficult (Heilman & Baltimore 1998 Nat. Med. 4 (4 Suppl.) 532-534). Although some success was reported in inducing nAb against laboratory HIV strains (Berman *et al*, 1990 Nature 345, 622-625; Fultz *et al*, 1992 Science 256, 1687-1690), it has been extremely difficult to neutralize primary isolates (Trkola *et al*, 1998 J. Virol. 72, 1876-1885; Haynes 1996 Lancet 34, 933-937). An explanation for the first 15 years of relative failure has been provided by the crystal structure of the core gp 120, which revealed multiple mechanisms by which HIV prevents efficient induction of nAb (Wyatt *et al*, 1998 Nature 393, 705-711; Kwong *et al*, 1998 Nature 393, 638-659). As a result of these difficulties, the emphasis of many vaccine designers has shifted to the induction of cell-mediated immune responses, which are mediated (predominantly) by cytotoxic T lymphocytes.

Cytotoxic T lymphocytes (CTL) are usually CD8⁺ cells and participate in an organism's defence in at least two different ways: they kill virus-infected cells; and they secrete a variety of cytokines and chemokines that directly or indirectly contribute to the suppression of virus replication. CTL-mediated protection after vaccination may depend on the levels of CTL present in the circulation and, perhaps, specifically for proteins expressed early (regulatory proteins) rather than late (structural proteins) in the replication cycle.

The induction and maintenance of CD8⁺ T cell responses require "help" provided by CD4⁺ T lymphocytes (helper T cells). In some HIV-infected individuals, high levels of HIV-specific helper response have been detected.

Identification of methods for induction of strong CD8⁺ T cell responses would provide tools for studying their role(s) in shaping the course of HIV infection and may stimulate progress towards an effective HIV vaccine. Previously, a prototype HIV vaccine was constructed as a string of partially overlapping epitopes recognised by murine, macaque and human CTL, which was delivered by vaccine vehicles that were safe and acceptable for use in humans, a DNA vector and modified vaccinia virus Ankara (MVA) vector (Hanke *et al*, 1998 Vaccine

16, 426-435; Hanke *et al*, 1998 J. Gen. Virol. 79, 83-90). In mice, the most potent protocol for induction of CTL was found to be DNA priming followed by MVA boosting (Hanke *et al*, 1998 Vaccine 16, 439-445; Schneider *et al*, 1998 Nat. Med. 4, 397-402) that is, priming mice with nucleic acid encoding the relevant polypeptide, followed by boosting the mice by inoculation with a modified vaccinia virus Ankara ("MVA") vector expressing the relevant epitopes.

WO 98/56919 discloses a "prime-boost" vaccination strategy, involving (i) priming with a composition comprising a source of one or more T cell epitopes of a target antigen, together with a pharmaceutically acceptable carrier, and (ii) boosting with a composition comprising a source of one or more T cell epitopes of the target antigen, including at least one T cell epitope that is the same as a T cell epitope of the priming composition.

The present invention aims, *inter alia*, to provide immunogens which may be useful in eliciting an HIV-specific response in humans. All documents and publications mentioned in this specification are incorporated herein by reference.

Summary of the Invention

In a first aspect the invention provides an immunogen in sterile form suitable for administration to a human subject, the immunogen comprising: at least a portion of the gag protein of HIV, said gag protein being from an HIV clade or having a consensus sequence for one or more HIV clades, and comprising at least parts of p17 and p24; and a synthetic polypeptide comprising a plurality of amino acid sequences, each sequence comprising a human CTL epitope of an HIV protein, and wherein a plurality of HIV proteins are represented in the synthetic polypeptide, said CTL epitopes being selected to stimulate an immune response to one or more HIV clades of interest.

For present purposes "sterile" refers to the general absence of viruses, bacteria, fungi, yeasts, chlamydia, mycoplasma, and spores of any of the foregoing (especially microbes pathogenic in humans). However, the immunogen may comprise one or more specific known microbial (e.g. viral or bacterial) vectors which serve to express the gag protein and/or synthetic polypeptide in a human subject. Such vectors are well known to those skilled in the art and

include, for example, viruses such as adenoviruses and pox viruses which are inherently non-pathogenic in humans or have been subjected to genetic manipulation or other modification to render them non-pathogenic in humans. Particularly preferred is vaccinia virus, especially modified vaccinia virus Ankara (MVA). Other specific preferred viral vectors include Semliki Forest Virus (SFV) and Sindbis virus (see Smerdon & Liljeström 2000, *Gene Ther. Regul.* 1, 1-31). Suitable bacterial vectors include BCG, and attenuated strains of *Salmonella* Spp. (especially "double aro" mutants of *Salmonella* which are being developed as vaccines for diarrhoeal diseases), and *Shigella* (see Shata *et al*, 2000 *Mol. Med. Today* 6, 66-71).

Other expression systems, which may be useful for producing the immunogen include a tobacco mosaic virus (TMV) expression vector (Palmer *et al*, 1999 *Arch. Virol.* 144, 1345-60) and NS1 tubules of bluetongue virus (Adler *et al*, 1998 *Med. Microbiol. Immunol. (Berl.)* 187, 91-96).

The term "synthetic" as used herein, is intended to refer to a polypeptide which is not, in its entirety, present in any naturally-occurring HIV isolate. The term "synthetic" is not intended to indicate that the polypeptide is necessarily synthesised by conventional chemical techniques of solid phase peptide synthesis. Whilst this represents one possibility, it is generally to be preferred that the polypeptide is synthesised by transcription and/or translation from an appropriate nucleic acid molecule encoding the synthetic polypeptide. Such methods of synthesis are well known to those skilled in the art and form no part of the invention.

It is preferred that the gag protein (or portion thereof) and the synthetic polypeptide are joined in some way on a single entity. For example, a viral vector may express both the gag protein and the synthetic polypeptide as separate components of the immunogen. Alternatively, both components of the immunogen may be covalently coupled or conjugated to each other or to a common carrier entity (such as a liposome, ISCOM or molecule). In preferred embodiments, both the gag protein and the synthetic polypeptide components of the immunogen are present in a single polypeptide or fusion protein. In such a fusion protein, the gag protein and synthetic polypeptide may be essentially the only components present. Alternatively the fusion protein may comprise other components which may

correspond to other HIV antigens (or portions thereof), or may be derived from other sources. Thus, in some embodiments, the polypeptide will comprise a portion of the gag protein substantially adjacent to the synthetic polypeptide (i.e. with fewer than 10 intervening amino acid residues). In other embodiments, the portion of the gag protein and the synthetic polypeptide may be separated by one or more intervening components (i.e. with 10 or more intervening amino acid residues), which will typically comprise one or more further HIV antigens.

It is also generally preferred that the immunogen does not contain the entire gag protein amino acid sequence. Typically between 55 and 95% of the gag protein will be present, preferably between 65 and 85%, most preferably about 75%.

The wild type HIV gag protein is known to consist of three portions termed p17, p24 and p15. These are synthesised in infected cells as a single polyprotein, with p17 at the N terminus. Normally, in HIV-infected cells, the N terminus of p17 is myristylated.

It is desirable that the gag portion of the immunogen will comprise at least part of p17 and p24, but it is generally preferred that the p17 component will be modified in some way to prevent myristylation. Conveniently this can be accomplished by reversing the order of the p17 and p24 components in the immunogen, such that p17 is no longer at a free N terminus and cannot therefore be myristylated. The inventors believe that this may improve the efficiency of presentation of peptides, derived from the immunogen, to a subject's immune system.

The gag protein component of the immunogen will generally comprise at least one T-helper cell, HLA Class II-restricted, peptide epitope and preferably comprise a plurality of such epitopes (preferably such that a number of different HLA Class II-restricted alleles are represented in the gag component). The gag protein component will also typically comprise one or more CTL HLA Class I-restricted peptide epitopes.

The synthetic polypeptide component of the immunogen will conveniently take the form of a string of CTL epitopes, each represented by, or contained within, a respective sequence of about 8-12 amino acids. Desirably at least some (preferably most) of the epitopes will be partially overlapping (such that one or more amino acids of one epitope will also be

contained within the sequence of an adjoining epitope). Some "non-epitopic" amino acid sequence may be present between neighbouring epitopes, but this is generally to be avoided.

Non-epitopic amino acid sequence between neighbouring epitopes is preferably less than 20 amino acid residues, more preferably less than 10 residues, and most preferably 1-5 amino acid residues. It will be apparent that such non-epitopic amino acid sequence may comprise linkers, spacers and the like which optimise the expression levels of the synthetic polypeptide or optimise its immunogenicity.

Thus in one extreme embodiment, all of the human CTL epitopes in the synthetic polypeptide may be overlapping and, at the other extreme, every epitope may be separated from its neighbours by at least some non-epitopic amino acid sequence. It is generally to be preferred that at least 50% of the human CTL epitopes are overlapping.

In addition to overlapping epitopes, the synthetic polypeptide may comprise at least some "adjacent epitopes". The term adjacent epitopes refers to epitopes which are not overlapping but which are not separated by any intervening non-epitopic amino acid sequence.

Thus in preferred embodiments the synthetic polypeptide comprises a mosaic pieced together from small (typically about 10-20 amino acid residue) fragments of different HIV proteins, which fragments will typically comprise one, two or three known adjacent and/or overlapping human CTL epitopes. Where a plurality of fragments are present in the synthetic polypeptide from the same HIV protein, the fragments will typically have been selected from discontinuous portions of the protein, so that it is unlikely that the synthetic polypeptide comprises a sequence corresponding to more than 20-25 consecutive amino acid residues of a particular HIV protein. Generally the synthetic polypeptide is designed so as essentially to omit those portions of HIV proteins not known to contain any human CTL epitopes.

A plurality of different HIV proteins will preferably be represented in the synthetic polypeptide. The synthetic polypeptide may contain epitopes present in any HIV antigen, but preferably will comprise at least one epitope present in one or more of the following: p24; pol; gp41; gp120; and nef. In one preferred embodiment, the synthetic polypeptide

comprises at least one CTL epitope present in each of the aforementioned HIV proteins. The synthetic polypeptide may additionally comprise at least one CTL epitope present in each of the following HIV proteins: vpr, vpu, vif (and especially) tat and rev.

It will be understood that the term "human CTL epitope" as used herein refers not to the origin of the protein from which the epitope derives, but indicates that the epitope is recognized and responded to by the CTL of at least a portion of the human population. Typically a human CTL epitope will be recognized by at least 0.01%, preferably 0.1%, and more preferably at least 1% of the world's human population.

In one particular embodiment, the immunogen specifically excludes any epitope from the env protein generally recognised by the human immune system. Most presently-used diagnostic tests are based on detection of an HIV env-specific immune response, so by excluding env components from the immunogen it is possible to distinguish between immune responses rising from infection with the virus and inoculation with the immunogen.

In one embodiment, the CTL epitopes present in the synthetic polypeptide are selected such that an immune response to HIV clade A will be generated. Preferably however, the synthetic polypeptide is large enough, and the CTL epitopes appropriately selected, such that an immune response which is cross-reactive against different HIV clades will be stimulated. This can conveniently be achieved by including one or more CTL epitopes which are conserved among different HIV clades. Several such epitopes are known to those skilled in the art (see Table 1 below).

In a preferred embodiment, the immunogen comprises at least one epitope (conveniently a CTL epitope) which is recognized by one or more laboratory test mammals, (e.g. mouse and/or monkey). Such an epitope can readily be incorporated within the synthetic polypeptide. Inclusion of an epitope of this sort allows for the quality, reproducibility and/or stability of different batches of the immunogen to be assayed in a potency assay using the laboratory test mammal (such as a mouse or macaque monkey). Examples of such epitopes include the amino acid sequence ACTPYDINQML (Seq. ID No. 1; containing a dominant epitope derived from simian immunodeficiency virus, SIV, gag p27, recognised by rhesus

macaque monkey CTLs) and RGPGRAFVTI (Seq. ID No. 2; an H-2D^d-restricted CTL mouse epitope derived from HIV env protein).

A list of some 23 human CTL epitopes suitable for inclusion in the synthetic polypeptide is shown in Table 1. The list is not exhaustive. A preferred immunogen will comprise at least 10 of such human CTL epitopes, more preferably at least 15, most preferably at least 20. In a particular embodiment each of the 23 human CTL epitopes listed in Table 1 will be represented in the synthetic polypeptide, optionally together with the macaque and murine epitopes also listed in Table 1.

Desirably the immunogen may additionally comprise a small tag or marker. Such a tag or marker should be as small as possible, in order to minimise the amount of extraneous material present in the immunogen. A convenient tag or marker allows for detection of expression and/or quantification of the amount of immunogen. Suitable tags include epitopes recognized by the monoclonal antibody Pk (Hanke *et al*, 1992 J. Gen. Virol. 73, 653-660).

The immunogen of the invention will conveniently be mixed with other substances in order to provide a vaccine composition. For example, a vaccine composition may additionally comprise one or more of the following: an adjuvant (e.g. alum), liposome, or immunostimulatory complex (ISCOM). A vaccine will also generally comprise a sterile diluent, excipient or carrier, such as a physiologically acceptable liquid (e.g. saline or phosphate-buffered saline solutions). The vaccine may be presented as a liquid or, more preferably, as a freeze-dried solid, which is suspended or dissolved in a physiologically acceptable liquid prior to administration.

Methods of administering the immunogen to a human subject will be apparent to those skilled in the art. Such methods include, in particular, intramuscular injection, subcutaneous injection, or delivery through the skin by needleless injection device. Alternatively, vaccines (especially those comprising bacterial vectors e.g. attenuated *Salmonella* or *Shigella* spp.) may be given orally, intranasally or by any other suitable route. A suitable dose of immunogen may typically be from 1 to 500 mg, typically from 10 to 100mg, depending on the size of the immunogen, the body mass of the recipient etc. A suitable dose can, if necessary, be ascertained by routine trial-and-error with different groups of subjects being

given different doses of immunogen, and the immune response of the subjects to the immunogen assayed in order to determine optimum dose size. The immune response of the subjects can be assayed by conventional immunological techniques (e.g. chromium release assay, using peripheral blood lymphocytes obtained from subjects' blood samples, acting on peptide-pulsed or virus-infected chromium-labelled target cells).

In a second aspect, the invention provides a nucleic acid molecule encoding a fusion protein, the fusion protein comprising an immunogen in accordance with the first aspect of the invention defined above.

The nucleic acid molecule is preferably in isolated, sterile form, suitable for administration to a human subject. The nucleic acid molecule is preferably "humanized" i.e. employs codons to code particular amino acids, which codons are frequently used in highly expressed human genes (Andre *et al*, 1998 J. Virol. 72, 1497-1503), instead of those codons used by HIV.

Desirably the nucleic acid molecule is contained within a vector, the sequence encoding the immunogen being operably linked to a promoter sequence active in human cells. Conveniently the promoter is a strong viral promoter such as the promoter from human cytomegalovirus (CMV). The vector will preferably also comprise an enhancer and polyadenylation signals, which are functional in human cells. Ideally, for maximum safety, the vector should not contain any origin of replication functional in human cells, to prevent undesirable replication of the vector.

The vector may be administered to the subject in isolation, as essentially 'naked' nucleic acid (preferably DNA), or else may be packaged within a delivery means, such as a virus, bacterium, liposome, or gold-coated particles and the like. One suitable delivery means is the modified vaccinia virus Ankara (MVA): the immunogen gene or open reading frame (ORF) may be inserted into MVA, for example, at the thymidine kinase locus.

It will be appreciated by those skilled in the art that a vector may comprise a nucleic acid encoding an immunogen (the nucleic acid therefore being in accordance with the second aspect of the invention), and that the vector may also possess, at its surface or internally, the polypeptide immunogen in accordance with the first aspect of the invention.

Whether administered as naked nucleic acid, or packaged within a delivery means, at least some of the administered nucleic acid enters the cells of the subject and is transcribed (if necessary) and translated, resulting in the *in situ* synthesis of the immunogen in the subject, who then develops an immune response to the immunogen.

As with the immunogen preparation *per se*, a nucleic acid encoding the immunogen may be administered to a human subject by any of a number of known routes e.g. subcutaneous or intramuscular injection, oral delivery, or delivery through the skin by means of needleless injector. Particular examples of administration by needleless injection device are disclosed in WO 97/34652 and WO 97/48485. A suitable dose of nucleic acid may be from 10 μ g to 10mg, typically from 100 μ g to 1mg, depending on the size of nucleic acid molecule, route of administration, body mass of the recipient etc. Again, routine trial and error (with the benefit of the present teaching) will enable those skilled in the art to determine an optimum dose of the nucleic acid.

Successful delivery of DNA to animal tissue has been achieved by cationic liposomes [Watanabe et al., Mol. Reprod. Dev. 38:268-274 (1994); Sharkey *et al*, WO 96/20013], direct injection of naked DNA into animal muscle tissue [Robinson et al., Vacc. 11:957-960 (1993); Hoffman et al., Vacc. 12:1529-1533; (1994); Xiang et al., Virol. 199:132-140 (1994); Webster et al., Vacc. 12:1495-1498 (1994); Davis et al., Vacc. 12:1503-1509 (1994); and Davis et al., Hum. Molec. Gen. 2:1847-1851 (1993)], and embryos [Naito et al., Mol. Reprod. Dev. 39:153-161 (1994); and Burdon et al., Mol. Reprod. Dev. 33:436-442 (1992)], or intradermal injection of DNA using "gene gun" technology [Johnston et al., Meth. Cell Biol. 43:353-365 (1994)].

For DNA-based vaccination, delivery by injection of naked plasmid DNA has shown potential in mouse models for inducing both humoral and cellular immune responses. However, in larger animals, using DNA delivery for vaccination has been hampered by requiring large amounts of DNA or inducing persistent expression of an antigen with the potential for developing tolerance to the antigen. Berglund reported a strategy for inducing or enhancing an immune response by injecting mice with plasmid DNA

containing an alphavirus DNA expression vector having a recombinant Semliki Forest Virus (SFV) replicon in a eukaryotic expression cassette [Berglund et al., *Nature Biotechnol.* 16:562-565 (1998)]. The eukaryotic expression cassette controlled expression of the primary nuclear transcription of the SFV replicon. This SFV replicon transcript, encoding the heterologous antigen, was transported to the cytoplasm and amplified by the self-encoded SFV replicase complex. The amplified RNA replicon lead to high level production of an mRNA encoding the heterologous antigen. Similar results were described by Polo and his group [Polo et al., *Nature Biotechnol.* 16:517-518 (1998); Hariharan et al., *J. Virol.* 72:950-958 (1998)]. Both groups found strong immune responses could be induced using small amounts of input plasmid DNA.

Alternatively, a method to deliver DNA to animals that overcomes the disadvantages of conventional delivery methods is by administering attenuated, invasive bacteria containing a bacterial DNA vector having a eukaryotic expression cassette encoding the gene to be expressed. For example, U.S. Patent No. 5,877,159 to Powell et al., describes live bacteria that can invade animal cells without establishing a productive infection or causing disease to thereby introduce a eukaryotic expression cassette encoding an antigen capable of being expressed by the animal cells.

In a third aspect the invention provides a method of stimulating an anti-HIV immune response in a human subject, the method comprising preparing an immunogen in accordance with the first aspect of the invention, or a nucleic acid molecule in accordance with the second aspect of the invention; and administering said immunogen or nucleic acid molecule to the subject.

Conveniently, the method comprises the administering both the immunogen and the nucleic acid molecule. In particular, the method preferably comprises one or more administrations of the nucleic acid molecule ("priming") followed at a suitable interval (e.g. 1 week to 4 months) by one or more administrations of the immunogen ("boosting"). Boosting may be performed, for example, by administering a replication-competent (e.g. attenuated virus or bacterium) or non-replicating vector comprising the immunogen and/or a nucleic acid

molecule encoding the immunogen. Preferably boosting is achieved by administering the immunogen as part of an MVA viral particle, which particle may advantageously comprise a nucleic acid encoding the immunogen.

In a preferred embodiment, performance of the method will result in the generation in the subject of a protective immune response such that, should the subject subsequently be exposed to HIV infection, the subject will not go on to develop the symptoms of AIDS associated with HIV infection.

In a fourth aspect the invention provides for use of an immunogen in accordance with the first aspect of the invention and/or a nucleic acid in accordance with the second aspect of the invention in the preparation of a medicament to prevent or treat HIV infection in a human subject.

In a fifth aspect, the invention provides a nucleic acid sequence encoding the amino acid sequence shown in Figure 8A. Conveniently the nucleic acid comprises or essentially consists of the nucleotide sequence shown in Figure 8B.

In a sixth aspect, the invention provides a polypeptide comprising the amino acid sequence shown in Figure 8A.

The nucleic acid of the fifth aspect and/or the polypeptide of the sixth aspect may be used in immunogen/vaccine compositions as described in relation to the other aspects of the invention, and such immunogens and vaccines are accordingly considered within the scope of the invention, and may comprise vectors etc (especially MVA) as aforesaid. The invention further provides, in a seventh aspect, a method of stimulating an anti-HIV immune response in a human subject, the method comprising administering to the subject a nucleic acid in accordance with the fifth aspect of the invention and/or a polypeptide in accordance with the sixth aspect of the invention. Finally, the invention provides for use of a nucleic acid in accordance with the fifth aspect of the invention and/or a polypeptide in accordance with the sixth aspect of the invention in the preparation of a medicament to prevent or treat HIV infection in a human subject.

The invention will now be further described by way of illustrative example and with reference to the accompanying drawings, wherein:

Figure 1A is a schematic representation of immunogens, HIV A, HIVTA and HIVAeT in accordance with the invention; and an immunogen PPA;

Figure 1B shows the amino acid sequence of the HIVA immunogen (Seq. ID No. 26);

Figure 2A shows the nucleotide sequence (Seq. ID No. 27) of a nucleic acid (termed "HIVA gene") encoding the HIV A immunogen;

Figure 2B is a schematic representation of the method used to construct the HIVA gene;

Figure 3 is a schematic representation of a DNA vector molecule (pTHr. HIVA) in accordance with the invention, and the method of its construction;

Figure 4 is a micrograph showing immunofluorescent detection of HIVA expression by mouse cells following transfection with pTHr. HIVA;

Figures 5A, B & C are graphs showing the results of chromium release assays using a splenocytes obtained from mice inoculated with a DNA molecule or an immunogen in accordance with the invention;

Figure 6A shows the amino acid sequence of the HIV TA immunogen (Seq. ID No. 28);

Figure 6B shows the nucleotide sequence (Seq. ID No. 29) of a nucleic acid molecule encoding the HIV TA immunogen;

Figure 7A shows the amino acid sequence (Seq. ID No. 30) of the tat polypeptide present in the HIVAeT immunogen;

Figure 7B shows the nucleotide sequence (Seq. ID No. 31) of a nucleic acid molecule encoding the HIVAeT immunogen;

Figure 8A shows the amino acid sequence (Seq. ID No. 32) of the PPA immunogen which is in accordance with the sixth aspect of the invention; and

Figure 8B shows the nucleotide sequence (Seq. ID No. 33) of a nucleic acid molecule (in accordance with the fifth aspect of the invention) encoding the PPA immunogen.

EXAMPLES

Example 1

This example relates to an immunogen for use in a vaccine focusing on the induction of cellular immune responses mediated by a concerted action of CD4⁺ helper and CD8⁺ effector T lymphocytes. The immunogen, designated HIVA (Hanke & McMichael Nat. Med. 6, 951-955), was designed for a phase III efficacy trial in Nairobi, Kenya. Figure 1A is a schematic representation of several immunogens, including HIVA. HIVA is derived from the sequences of HIV-1 clade A, the predominant HIV clade in Nairobi and consists of about 73% of the gag protein fused to a string of 25 partially overlapping CTL epitopes. The gag domain of HIVA contains p24 and p17 in an order reversed to the viral gag p17-p24-p15 polyprotein. This rearrangement prevents myristylation of the N-terminus of p17, which could direct the recombinant protein to the cell membrane, thus preventing efficient degradation into peptides necessary for the major histocompatibility complex (MHC) class I presentation.

Figure 1B shows the amino acid sequence (Seq. ID No. 26) of the HIVA immunogen. Amino acids corresponding to the restriction endonuclease sites used to assemble the gene are shown in bold (GS-corresponds to *Bam* HI, GT corresponds to *Kpn*I and EF corresponds to *Eco*RI).

The amino acid sequence of the gag domain was derived from the protein database consensus sequence of HIV-1 clade A (Korber *et al*, "Human retroviruses and AIDS; a compilation and analysis of nucleic acid and amino acid sequences" 1997). In the absence of

available Kenyan strain sequences, regions without a strong amino acid clade A preference were biased towards Ugandan isolates. The HIV-1 gag protein contained not only important MHC class I-, but also class II-restricted epitopes which stimulate CD4⁺ T helper cells.

The C-terminus of the HIVA protein was designed as a multi-CTL epitope synthetic polypeptide. The CTL epitopes included in HIVA were recognised by CTL in patients infected with HIV-1 clade A strains circulating in Kenya, were 8- to 10-amino acids long, and originate from the gag, pol, nef or env proteins (Rowland-Jones *et al*, 1998 J. Clin. Invest. 102, 1758-1765; Dorrell *et al*, 1999 J. Virol. 73, 1708-1714). Many of these epitopes are immunodominant and relatively conserved among other HIV-1 clades (Table 1) (and therefore should be able to elicit an immune response which cross-reacts with HIV viruses of clades other than clade A). They are presented by seventeen different HLA alleles, which include both frequent African alleles as well as alleles common in most ethnic populations. It has been estimated that optimally selected epitopes presented by the nine commonest HLA alleles could cover the general population irrespective of ethnic descent (Sydney *et al*, 1996 Immunol. Today 17, 261-266). Thus, given that majority of HIV-infected donors make good CTL responses to gag p17/p24, each vaccinee should have the potential to respond to at least two or three CTL epitopes present in the HIVA protein.

The HIVA synthetic polypeptide also comprised SIV gag and HIV env epitopes recognised by macaque and murine CTL, respectively, so that the quality, reproducibility and stability of the clinical batches could easily be assessed in a mouse (or macaque if necessary) potency assay. A monoclonal antibody epitope Pk (Hanke *et al*, 1992 J. Gen. Virol. 73, 653-660) was added to the C-terminus of HIVA for easy detection of the full-size protein and estimation of the level of expression. There is no reason to believe that the three non-HLA epitopes represent a health hazard for the vaccinated individuals.

Table 1. CD8+ cell epitopes included in the HIVA synthetic polypeptide polyepitope region.

Epitope ^a	MHC class I restriction	Origin	HIV clade ^b
ALKHRAYEL	HLA-A*0201	nef	a
PPIPVGIEY	HLA-B35	p24	a/B/c/D/F/G
GEIYKRWII	HLA-B8	p24	a/B/c/D/F/G
KRWIILGLNK	HLA-B*2705	p24	A/B/C/D/F/G/H
FRDYVDRFYK	HLA-B18	p24	BD(A=C/F/G/H) ^c
RDYVDRFYKTL	HLA-B44	P24	B/D(A=C/F/G/H) ^c
DRFYKTLRA	HLA-B14	p24	B/D(A=C/F/G/H)
AIFQSSMTK	HLA-A*0301, A11, A33	pol	a/B/c/D/G/H
ITLWQRPLV	HLA-A*6802	pol	a/b/C/D/F/G/H
ERYLKDQQL	HLA-B14	gp41	a/b/C/D
YLKDQQLL	HLA-A24, B8	gp41	a/b/C/D
TVYYGVPVWK	HLA-A*0301	gp120	A/B/C/D/g
RPQVPLRPMTY	HLA-B51	nef	A/b/D/E/F/G
QVPLRPMTYK	HLA-A*0301, A11	nef	A/b/D/E/F/G
VPLRPMTY	HLA-B35	nef	A/b/D/E/F/G
AVDLSHFLK	HLA-A11	nef	a/B/d/f
DLSHFLKEK	HLA-A*0301	nef	A/B/D/F
FLKEKGGL	HLA-B8	nef	A/B/C/D/E/F/G
ILKEPVHGV	HLA-A*0201	pol	A/B/C/D/G
ILKEPVHGVY	HLA-Bw62	pol	A/B/D
HPDIVTYQY	HLA-B35	pol	a
VIYQYMDDL	HLA-A*0201	pol	A/B/C/D/F/G/H
TPGPGVRYPL	HLA-B7	nef	b/c
ACTPYDINQML ^d	Mamu-A*01	p27	SIV
RGPGRAFVTI ^e	H-2D ^d	env	HIV

- a* - Epitopes are listed (Seq. ID Nos. 3-25, 1, 2) in the order in which they appear in the polypeptide.
- b* - A particular epitope sequence is present in about 50% (small letter) or 90% (capital letter) of sequenced HIV clade isolates.
- c* - '=' indicates that the epitopes are present in the N-terminal clade A gag domain.
- d* - A dominant epitope derived from SIV gag p27 flanked by Ala and Leu at its N- and C- termini, respectively, recognised by rhesus macaque CTL, which can be used for potency studies in rhesus macaques.
- e* - A CTL epitope presented by a murine MHC class I used for the mouse potency assay.

Since it was intended to adopt a "prime/boost" protocol, in which priming was achieved by administering nucleic acid, it was desirable to design the sequence of the nucleic acid encoding the HIVA immunogen in order to increase the expression of HIVA in human cells. Firstly, to ensure an efficient initiation of translation from the first methionine codon, the HIVA open reading frame (ORF) was preceded by a 12-nucleotide-long Kozak consensus sequence (Kozak 1987 Nucl. Acids Res. 15, 8125-8148). Secondly, the translation of the resulting mRNA was optimised by substituting most of the HIV-1-derived codons with frequently used codons in highly expressed human genes (Andre *et al*, 1998, cited above). The HIVA ORF is a part of a 1,608-base pair-long double-stranded DNA *HindIII-XbaI* fragment. (Figure 2A shows the nucleotide sequence of the *HindIII-XbaI* insert containing the HIVA ORF; Seq. ID No. 27) In Figure 2A, the endonuclease sites used to assemble the partial PCR products are included (nucleotides 1-6 = *HindIII*; 319-324 = *Bam* HI; 712-717 = *Kpn* I; 1135-1140 = *EcoRI*; and 1603-1608 = *XbaI*).

Plasmid DNA was prepared and treated using standard protocols (Sambrook *et al*, "Molecular Cloning. A Laboratory Manual" 2nd Edition; Cold Spring Harbor). The HIVA gene was constructed (as indicated in Figure 2B) *in vitro* in four parts. Each part was prepared by assembly from overlapping positive- and negative-strand oligodeoxynucleotides of 70-90 bases in length. The synthetic oligodeoxynucleotides were

purified using the EconoPure™ Kit (Perkin Elmer) according to the manufacturer's instructions, annealed and ligated, followed by PCR assembly. The PCR products were gel-purified after each step until fragments with the expected, unique terminal restriction endonuclease sites were obtained. These four products were cloned and sequenced, and ligated together to generate the complete HIVA gene. The HIVA gene was then inserted into the pTHr construct and MVA vaccine vector, as described below.

The pTHr vector. A vector pTHr for a direct gene transfer was designed with the aim to minimise the number of functional elements and therefore the amount of DNA required to be administered.

The construction of pTH was described previously (Hanke *et al*, 1998 Vaccine 16, 426-435). It contains an expression efficient enhancer/promoter/intron A cassette of the human cytomegalovirus (hCMV) strain AD169 genome (Whittle *et al*, 1987 Protein Eng. 1, 499-505; Bebbington 1991 Methods 2, 136-145). The promoter region is followed by the pRc/CMV (Invitrogen)-derived polylinker and polyadenylation signal of the bovine growth hormone gene. The β -lactamase gene conferring ampicillin resistance to transformed bacteria and prokaryotic origin of double-stranded DNA replication ColE1 are both derived from plasmid pUC19. pTH does not contain an origin for replication in mammalian cells. After insertion of the HIVA DNA into the pTH polylinker, the β -lactamase gene fragment between the *Mlu*I and *Dra*I sites was removed and the resulting construct pTHr.HIVA (Fig. 3) was propagated in bacteria using the repressor-titration system developed by Cobra Pharmaceuticals Ltd. (Keele, UK), which selects plasmid-carrying bacteria without the need for the presence of an antibiotic-resistance gene on the plasmid (Williams *et al*, 1998 Nucl. Acids Res. 26, 2120-2124). Therefore, DNA vaccination does not introduce into the human vaccinee large numbers of copies of an antibiotic resistance gene. Construction of pTHr. HIVA is illustrated schematically in Figure 3 (CMV e/p/i = human CMV enhancer/promoter/intron A cassette; BGHpA = bovine growth hormone polyadenylation signal; ColE1 = origin of dsDNA replication in bacteria; arrow head symbols denote repressor-binding sequences).

293T cells (and chicken embryo fibroblasts [CEF] were maintained in Dulbecco's modified Eagle's medium (DMEM; Gibco) supplemented with 10% foetal bovine serum (FBS; Gibco); 2mM L-glutamine and penicillin/streptomycin. Cells were cultured in a humidified incubator in 5% CO₂ at 37°C. 293T cells were transiently transfected with pTHr. HIVA using the DEAE-dextran-chloroquine method (Hanke *et al*, 1998 Vaccine 16, 426-435). Briefly, 2.5×10^5 293T cells were grown on coverslips in 6-well tissue culture plates overnight. The following day, cells were transfected with 5 g per well of DNA. After 48 hours, the transfected cells were fixed, their membranes were permeabilized and the SV5-P-k mAb followed by anti-murine FITC-conjugated antibodies were used to detect the expressed recombinant proteins.

A micrograph illustrating specimen results is shown in Figure 4. The Figure shows three transfected cells and one background untransfected cell (top left).

Boosting was to be achieved by administering an MVA vector expressing the HIVA immunogen. MVA is an attenuated vaccinia virus safe for clinical application, which has almost lost its ability to replicate in human cells (Mayr *et al*, 1975 Infection 105, 6-14). The use of MVA as a vaccine vehicle and its features which make it an attractive choice among the attenuated poxvirus vectors (see, e.g. Sutter *et al*, 1994 Vaccine 12, 1032-1040; and Moss *et al*, 1996 Adv. Exp. Med. Biol. 397, 7-13) have been described extensively.

The HIVA gene was ligated into plasmid pSC11, which directed the gene into the thymidine kinase locus of the parental MVA (Carroll & Moss 1995 Biotechniques 19, 352-356). Bulk stocks of the recombinant MVA were grown on primary CEF obtained from the eggs of a specific pathogen-free flock. MVA was purified by centrifugation of cytoplasmic extracts through a 36% (w/v) sucrose cushion in a Beckman SW28 rotor at 13,500 rpm for 80 minutes. Taking advantage of the co-inserted β -galactosidase gene, the virus stock titres were determined from the number of blue plaques after incubation of the infected cell monolayers with the appropriate substrate.

Vaccine potency assay. The potencies of the DNA and MVA vectors were tested in groups of Balb/c mice taking advantage of the presence of the H-2D^d-restricted epitope (Takahashi *et al*, 1993 Int. Immunol. 5, 849-857). For the pTHr.HIVA DNA vaccine, mice were either needle-injected intramuscularly with 100 µg of DNA or immunised twice 3 weeks apart intradermally with total of 2 µg of DNA using the Dermal XR gene delivery device of PowderJect Vaccines Inc. (Madison, WI, USA). The mice were sacrificed 10 or 21 days after the last immunisation. Spleens from the immunized mice were removed and pressed individually through a cell strainer (Falcon) using a 2-ml syringe rubber plunger. The splenocytes were washed and divided into two halves. One half was frozen for tetramer analysis and the second half was suspended in 5ml of Lymphocyte medium (R10, 20 mM HEPES and 15 mM β-mercaptoethanol) and restimulated *in vitro* by incubation with 2 µg/ml of the RGPGRFVFI (Seq. ID No. 2) peptide in an humidified incubator in 5% CO₂ at 37°C for 5 days.

The effector cells were 2-fold diluted in U-bottom wells (96-well plate; Costar) starting with 100:1 effector to target ratio. Five thousand ⁵¹Cr-labeled P815 target cells in a medium without or supplemented with 10⁻⁶ M peptide was then added to the effectors and incubated at 37°C for 5 hours. Spontaneous and total chromium releases were estimated from wells, in which the target cells were kept in a medium alone or with 5% Triton X-100, respectively. The percentage specific lysis was calculated as [(sample release-spontaneous release)/(total release-spontaneous release)] x 100. The spontaneous release was lower than 5% of the total c.p.m.

The results of typical assays are shown in Figures 5A-C, which are graphs of % specific lysis against Effector: Target ratio. Figure 5A shows results for splenocytes obtained from mice immunised once with 100 µg of pTHr.HIVA DNA. Figure 5B shows results from mice immunised twice intradermally with pTHr. HIVA DNA. Figure 5C shows results obtained from mice immunised intramuscularly with 10⁷ pfu of MVA.HIVA. In each of Figures 5A-5C, each line represents a single mouse. Lysis of peptide-pulsed targets is denoted by filled symbols, unpulsed targets by open symbols.

Both modes of the pTHr.HIVA vector delivery were highly immunogenic and induced high cytolytic activities in all immunised animals (see Figs. 5A & B). Similarly, a single needle intramuscular injection of 10^7 plaque-forming units of MVA.HIVA elicited in all vaccinees strong peptide-specific lytic activities measured after a culture restimulation (Fig. 5C).

Example 2

Further nucleic acid constructs were prepared which encoded polyprotein immunogens based on HIVA, but comprising further HIV antigen components. These further constructs and their encoded immunogens were termed HIVTA and HIVAeT, as shown in Figure 1A. Also shown is a construct/immunogen termed PPA. The relevant DNA/amino acid sequences are shown in Figures 6 A/B-8 A/B respectively, although Figure 7A shows only that portion of amino acid sequence in HIVA eT (attributable to tat) which is additional to that in HIVTA. (Note that the sequence of the *HindIII* and *XbaI* sites at the 5' and 3' ends of the DNA sequences are not shown in Figures 6B, 7B and 8B). The constructs were prepared in a manner similar to that described above for HIVA.

HIVTA and HIVAeT share the same design rationale with HIVA, but additionally include the HIV-1 clade A tat sequence, expressed either as part of a fusion protein with gag and the polyepitopic synthetic polypeptide (in the case of HIVTA, the tat sequence being positioned between gag and the synthetic polypeptide), or else being present on the same construct but expressed as a separate polypeptide (in the case of HIVAeT), by virtue of the inclusion of an internal ribosome entry site (IRES).

Each of the nucleic acid molecules may be used to immunise subjects, in a manner similar to that described above in relation to HIVA DNA. Equally, the molecules may be introduced into appropriate vectors, especially MVA, again as described above in Example 1, and the resulting vector used to immunise subjects.

The significance of the genetic diversity among individual HIV isolates and its implication for vaccine design have been long debated. The predominant HIV-1 clade in Europe and

North America is clade B, which is also the most studied one. In central and Eastern Africa, the predominant circulating HIV-1 strain is clade A, while clade C is dominating Southern Africa, India and China. Generally, a clade-specific vaccine design requires a more careful consideration for the induction of nAb than for CTL. Although there are some important inter-clade differences in CTL epitopes, many epitopes are conserved across clades partially due to structure/function constraints. However, to facilitate the interpretation of efficacy studies, vaccines should attempt to match the local strains prevalent in the trial population with the view that any successful approaches can be adapted for other clades if cross-protection is not achieved.

Claims

1. An immunogen in sterile form suitable for administration to a human subject, the immunogen comprising: at least a portion of the gag protein of HIV, said gag protein being from an HIV clade or having a consensus sequence for one or more HIV clades, and comprising at least parts of p17 and p24; and a synthetic polypeptide comprising a plurality of amino acid sequences, each sequence comprising a human CTL epitope of an HIV protein, and wherein a plurality of HIV proteins are represented in the synthetic polypeptide, said CTL epitopes being selected to stimulate an immune response to one or more HIV clades of interest.
2. An immunogen according to claim 1, in which the said at least portion of gag protein and the synthetic polypeptide are present as a fusion protein.
3. An immunogen according to claim 1 or 2, in which at 55-95% of the amino acid sequence of gag protein is present in the portion of gag.
4. An immunogen according to any one of claims 1, 2 or 3, in which 65-85% of the amino acid sequence of gag protein is present in the portion of gag.
5. An immunogen according to any one of the preceding claims, in which said at least part of p17 is modified to prevent N-terminal myristylation.
6. An immunogen according to any one of the preceding claims, in which at least some of the human CTL epitopes present in the synthetic polypeptide are overlapping.
7. An immunogen according to claim 6, in which at least 50% of the human CTL epitopes present in the synthetic polypeptide are overlapping.
8. An immunogen according to any one of the preceding claims, in which at least some of the human CTL epitopes present in the synthetic polypeptide are adjacent.

9. An immunogen in according to any one of the preceding claims, in which at least some of the human CTL epitopes are separated by non-epitopic amino acid sequence.
10. An immunogen according to any one of the preceding claims, comprising at least one epitope which is recognised by one or more laboratory test mammals.
11. An immunogen according to claim 10, wherein said epitope recognised by one or more laboratory test mammals is a CTL epitope.
12. An immunogen according to any one of the preceding claims, in which the synthetic polypeptide comprises at least 10 of the human CTL epitopes identified in Table 1.
13. An immunogen according to any one of the preceding claims, in which the synthetic polypeptide comprises at least 15 of the human CTL epitopes identified in Table 1.
14. An immunogen according to any one of the preceding claims, in which the synthetic polypeptide comprises at least 20 of the human CTL epitopes identified in Table 1.
15. An immunogen according to any one of the preceding claims, in which the synthetic polypeptide comprises all 23 of the human CTL epitopes identified in Table 1.
16. An HIV immunogen for a human subject comprising at least a portion of a gag protein, said gag protein being from an HIV clade of interest or having a consensus sequence for an HIV clade of interest, and wherein said portion comprises at least parts of p17 and p24 and is modified to prevent N-terminal myristylation; and a synthetic polypeptide comprising a plurality of human CTL epitopes from a plurality of different HIV proteins, each CTL epitope having from 8 to 12 amino acids, and said CTL epitopes being selected to stimulate an immune response to the clade of interest.

17. An HIV immunogen for a human subject comprising at least a portion of a gag protein, said gag protein being from an HIV clade of interest or having a consensus sequence for an HIV clade of interest, and wherein said portion comprises at least parts of p17 and p24 and is modified to prevent N-terminal myristylation; and a synthetic polypeptide consisting essentially of a plurality of human CTL epitopes from a plurality of different HIV proteins, each CTL epitope having from 8 to 12 amino acids, and said CTL epitopes being selected to stimulate an immune response to said clade of interest; and optionally, said synthetic polypeptide having at least one immunogenic epitope recognised by a laboratory animal.
18. An immunogen according to any one of the preceding claims, comprising the amino acid sequence shown in Figure 1B.
19. An immunogen according to claim 18, consisting essentially of the amino acid sequence shown in Figure 1B.
20. An immunogen according to any one of the preceding claims, wherein said clade of interest is selected from the group consisting of HIV clade A, B, C, D, E, F, G and H.
21. An immunogen according to claim 20, wherein said clade of interest is HIV clade A.
22. An immunogen according to claim 20, wherein said clade of interest is HIV clade B.
23. An immunogen according to claim 20, wherein said clade of interest is HIV clade C.
24. An immunogen according to any one of the preceding claims, wherein said human CTL epitopes include at least one epitope from each of the HIV proteins nef, p24, p17, pol, gp41 and gp120.

25. An immunogen according to claim 24, wherein wherein said human CTL epitopes include at least one epitope from each of the HIV proteins nef, p24, p17, pol, gp41, gp120, tat, and rev.
26. An immunogen according to claim 24 wherein said human CTL epitopes include at least one epitope from each of the HIV proteins nef, p24, p17, pol, gp41, gp120, tat, rev, vpr, vpu and vif.
27. A nucleic acid molecule encoding an immunogen in accordance with any one of the preceding claims.
28. A nucleic acid according to claim 27, wherein the immunogen is encoded as a fusion protein.
29. A nucleic acid molecule according to claim 27 or 28, comprising the nucleotide sequence shown in Figure 2A.
30. A vector comprising a nucleic acid molecule in accordance with any one of claims 27, 28 or 29.
31. A particulate vector according to claim 30, further comprising an immunogen in accordance with any one of claims 1-26.
32. A method of stimulating an anti-HIV immune response in a human subject, the method comprising the steps of: preparing an immunogen in accordance with any one of claims 1-26 and/or a nucleic acid molecule in accordance with any one of claims 27-29; and administering the immunogen and/or the nucleic acid molecule to the subject.

33. A method according to claim 32, wherein the subject is primed by one or more administrations of the nucleic acid molecule; and subsequently boosted by one or more administrations of the immunogen.
34. Use of an immunogen in accordance with any one of claims 1-26 and/or a nucleic acid molecule in accordance with any one of claims 27-29, in the preparation of a medicament to prevent or treat HIV infection in a human subject.
35. A method of stimulating an anti-HIV immune response in a human subject which comprises administering one or more times an amount of a nucleic acid encoding an immunogen in accordance with any one of claims 1-26 to said subject sufficient to prime an immune response to said immunogen, and administering one or more times a modified vaccinia virus Ankara (MVA) particle encoding and/or containing an immunogen in accordance with any one of claims 1-26 to said subject in an amount sufficient to boost the immune response to common portions of said immunogens.
36. A bacterium comprising an immunogen in accordance with any one of claims 1-26, or comprising a nucleic acid molecule in accordance with any one of claims 27-29.
37. A bacterium according to claim 36 which is an attenuated pathogen suitable for administration to a human subject.
38. A nucleic acid molecule encoding a polypeptide comprising the amino acid sequence shown in Figure 8A.
39. A nucleic acid molecule encoding a polypeptide consisting essentially of the amino acid sequence shown in Figure 8A.
40. A nucleic acid molecule according to claim 38 or 39, comprising the nucleotide sequence shown in Figure 8B.

41. A polypeptide comprising the amino acid sequence shown in Figure 8A.
42. A polypeptide consisting essentially of the amino acid sequence shown in Figure 8A.
43. A particulate vector comprising the nucleic acid sequence of claim 38 or 39 and/or the polypeptide of claim 41 or 42.
44. A method of stimulating an anti-HIV immune response in a human subject, comprising the steps of: preparing a nucleic acid molecule in accordance with claim 38 or 39 and/or a polypeptide in accordance with claim 41 or 42, and administering the nucleic acid molecule and/or the polypeptide to the subject.
45. Use of a nucleic acid molecule in accordance with claim 38 or 39 and/or a polypeptide in accordance with claim 41 or 42, in the preparation of a medicament to prevent or treat HIV infection in a human subject.
46. An immunogen substantially as hereinbefore described and with reference to the accompanying drawings.
47. A method of stimulating an anti-HIV immune response in a human subject substantially as hereinbefore described and with reference to the accompanying drawings.

1/11

Fig.1A.




Fig.1B.

1	MPIVQNAQQG	MHQALSPRTL	NAWVKVIEEK	AFSPEVIPMF	SALSEGATPQ	50
51	DLNMLNIVG	GHQAAMQMLK	DTINEEAAEW	DRLHPVHAGP	IPPGQMRPR	100
101	GSDIAGTTST	LQEQIGWMTS	NPPIPVGDIY	KRWIILGLNK	IVRMYSFVSI	150
151	LDIRQGPKEP	FRDYVDRFFK	TLRAEQATQE	VKNWMTETLL	VQNANPDCKS	200
201	ILRALGPGAT	LEEMMTACQG	VGGPGHKARV	LGTGARASVL	SGGKLDAWEK	250
251	IRLRPGGKKK	YRLKHLVWAS	RELRFALNP	SLLETAEGCQ	QIMEQLQSAL	300
301	KTSEELKSLF	NTVATLYCVH	QRIDVKDTKE	ALDKIEEIQN	KSKQKTQQAA	350
351	ADTQSSSKVS	QNYALKHRAV	ELEFPPIPVG	EIYKRWIIFR	DYVDRFYKTL	400
401	RAIFQSSMTK	ITLWQRPLVE	RYLKDQQLLT	VYYGVPVWKR	PQVPLRPMTY	450
451	KAVDLSHFLK	EKGGLILKEP	VHGVYHPDIV	IYQYMDDLTP	GPGVRYPLAC	500
501	TPYDINQMLR	GPGRAFVTIP	NPLLGLD			527

2/11


Fig.2A.

HIVA DNA

AAGCTTCCCGCCGCCACCATGCCCATCGTGCAGAACGCCAGGGCCAGATGCACCAGGCC
CTGTCCCCCGCACCCCTGAACGCCTGGGTGAAGGTGATCGAGGAGAAGGCCTTCTCCCC
GAGGTGATCCCCATGTTCTCCGCCCTGTCCGAGGGCGCCACCCCCAGGACCTGAACATG
ATGCTGAACATCGTGGGCGGCCACCAGGCCGCCATGCAGATGCTGAAGGACACCATCAAC
GAGGAGGCCGCCGAGTGGGACCGCCTGCACCCCGTGCACGCCGGCCCCATCCCCCGGC
CAGATGCGCGAGCCCCGCGGATCCGACATCGCCGGCACCACCTCCACCCTGCAGGAGCAG
ATCGGCTGGATGACCTCCAACCCCCCATCCCCGTGGGCGACATCTACAAGCGCTGGATC
ATCCTGGGCCTGAACAAGATCGTACGCATGTACTCCCCCGTGTCCATCCTGGACATCCGC
CAGGGCCCCAAGGAGCCCTTCCGCGACTACGTGGACCGCTTCTTCAAGACCCTGCGCGCC
GAGCAGGCCACCCAGGAGGTGAAGAACTGGATGACCGAGACCCTGCTGGTGCAGAACGCC
AACCCCGACTGCAAGTCCATCCTGCGCGCCCTGGGCCCCGCGGCCACCCTGGAGGAGATG
ATGACCGCCTGCCAGGGCGTGGGCGGCCCCGGCCACAAGGCCCGCGTGTGGGTACCGGC
GCCCCGCGCCTCCGTGCTGTCCGCGGCAAGCTGGACGCCTGGGAGAAGATCCGCCTGCGC
CCCGCGGCAAGAAGAAGTACCGCCTGAAGCACCTGGTGTGGGCCTCCCGCGAGCTGGAG
CGCTTCGCCCTGAACCCCTCCCTGCTGGAGACCGCCGAGGGCTGCCAGCAGATCATGGAG
CAGCTGCAGTCCGCCCTGAAGACCTCCGAGGAGCTGAAGTCCCTGTTCAACACCGTGGCC
ACCCTGTACTGCGTGCACCAGCGCATCGACGTGAAGGACACCAAGGAGGCCCTGGACAAG
ATCGAGGAGATCCAGAACAAGTCCAAGCAGAAGACCCAGCAGGCCGCCGCCGACACCCAG
TCCTCCTCCAAGGTGTCCCAGAACTACGCCCTGAAGCACCGCGCCTACGAGCTGGAATTC
CCTCCAATTCTGTGCGGGAGATTTATAAACGGTGGATCATTTTTAGGGATTATGTGAT
AGGTTTTATAAAACGCTCAGGGCCATCTTCCAGTCTCCATGACCAAGATCACCTGTGG
CAGCGCCCCCTGGTGGAGCGCTACCTGAAGGACCAGCAGCTGCTGACCGTGTACTACGGC
GTGCCCCGTGTGGAAGCGCCCCCAGGTGCCCTGCGCCCCATGACCTACAAGGCCGTGGAC
CTGTCCCACCTTCTGAAGGAGAAGGGCGGCCTGATCCTGAAGGAGCCCGTGCACGGCGTG
TACCACCCCGACATCGTGATCTACCAGTACATGGACGACCTGACCCCGGCCCGCGCTG
CGCTACCCCTGGCCTGCACCCCTACGACATCAACCAGATGCTGCGCGGCCCGGCCGC
GCCTTCGTGACCATCCCCAACCCCTGCTGGGCCTGGACTGATCTAGA


3/11

Fig.2B.




4/11

Fig.3.



5/11

Fig.4.



6/11

Fig.5A.




Fig.5B.





Fig.5C.



7/11

Fig.6B.

HIVTA DNA

CCCC GCCC ACCATG CCCATCGTGCAGAACGCC CAGGGCCAGATGCACCAGGCCCTGTCC
 CCCC GCACCCTGAACGCCTGGGTGAAGGTGATCGAGGAGAAGGCCTTCTCCCCGAGGTG
 ATCCCCATGTTCTCCGCCCTGTCCGAGGGCGCCACCCCCCAGGACCTGAACATGATGCTG
 AACATCGTGGGCGGCCACCAGGCCGCCATGCAGATGCTGAAGGACACCATCAACGAGGAG
 GCCGCCGAGTGGGACCGCCTGCACCCCGTGCACGCCGGCCCCATCCCCCCCCGCCAGATG
 CGCGAGCCCCGCGGATCCGACATCGCCGGCACCACCTCCACCCTGCAGGAGCAGATCGGC
 TGGATGACCTCCAACCCCCCATCCCCGTGGGCGACATCTACAAGCGCTGGATCATCCTG
 GGCCTGAACAAGATCGTGCAGATGTACTCCCCCGTGTCCATCCTGGACATCCGCCAGGGC
 CCCAAGGAGCCCTTCCGCGACTACGTGGACCGCTTCTTCAAGACCTGCGCGCCGAGCAG
 GCCACCCAGGAGGTGAAGAACTGGATGACCGAGACCCTGCTGGTGCAGAACGCCAACCCC
 GACTGCAAGTCCATCCTGCGCGCCCTGGGCCCCGCGCCACCCTGGAGGAGATGATGACC
 GCCTGCCAGGGCGTGGGCGGCCCGGCCACAAGGCCCGCGTGTGGGTACCGGCGCCCGC
 GCCTCCGTGCTGTCCGCGGCAAGCTGGACGCCTGGGAGAAGATCCGCCTGCGCCCCGGC
 GGCAAGAAGAAGTACCGCCTGAAGCACCTGGTGTGGGCCTCCCGCGAGCTGGAGCGCTTC
 GCCCTGAACCCCTCCCTGCTGGAGACCGCCGAGGGCTGCCAGCAGATCATGGAGCAGCTG
 CAGTCCGCCCTGAAGACCTCCGAGGAGCTGAAGTCCCTGTTCAACACCGTGGCCACCCTG
 TACTGCGTGCACCAGCGCATCGACGTGAAGGACACCAAGGAGGCCCTGGACAAGATCGAG
 GAGATCCAGAACAAAGTCCAAGCAGAAGACCCAGCAGGCCGCCGCCGACACCCAGTCTTCC
 TCCAAGGTGTCCAGAACTACGCCCTGAAGCACC CGCCTACGAGCTGGAATTCATGGCC
 ACAACCATGGACCCCGTGGACCCCAACCTGGAGCCCTGGAACCACCCCGGCTCCAGCCC
 ACCACCCCGGCTCCAAGTGCTACTGCAAGGTGTGCTGCTACCACTGCCCCGTGTGCTTC
 CTGAACAAGGGCCTGGGCATCTCCTACGGCCGCAAGAAGCGCCGCCAGCGCCGCGGCACC
 CCCCAGTCCAACAAGGACCACCAGAACCCCATCCCCAAGCAGCCCATCCCCAGACCCAG
 GGCATCTCCACCGGCTCCAAGGAGTCCAAGAAGAAGGTGGAGTCCAAGACCGAGACCGAC
 CCGGAGGAATTCCTCCAATTCCTGTGCGGGGAGATTTATAAACGGTGGATCATTTTATAGG
 GATTATGTCGATAGGTTTTATAAAACGCTCAGGGCCATCTTCCAGTCTCCATGACCAAG
 ATCACCTGTGGCAGCGCCCCCTGGTGGAGCGCTACCTGAAGGACCAGCAGCTGCTGACC
 GTGTACTACGGCGTGCCCGTGTGGAAGCGCCCCCAGGTGCCCCCTGCGCCCCATGACCTAC
 AAGGCCGTGGACCTGTCCCACTTCTGAAGGAGAAGGGCGGCCTGATCCTGAAGGAGCCC
 GTGCACGGCGTGTACCACCCCGACATCGTATCTACCAGTACATGGACGACCTGACCCCC
 GGCCCCGGCGTGCGCTACCCCTGGCCTGCACCCCTACGACATCAACCAGATGCTGCGC
 GGCCCCGGCCGCGCCTTCGTGACCATCCCCAACCCCTGCTGGGCCTGGACTGA

Fig.6A.

HIVTA protein

MPIVQNAQQQM HQALSPRTLNAWVKVIEEKAFSPEVIPMFSA LSEGATPQDLNM
 MLNIVGGHQAAMQMLKDTN EEA EWDR LHPVHAGPIPPGQMREPRGSDIAGT
 TSTLQEQIGWMTSNPPIPVGD IYKRWIILGLNKIVRMYS PVSILDIRQGPKEPFRDY
 VDRFFKTLRAEQATQEVKNWMTETLLVQNANPDCKSILRALGPGATLEEMMTA
 CQGVGGPGHKARVLGTGARASVLSGGKLD AWEKIRLRPGGKKKYRLKHLVWA
 SRELERFALNPSLLETAEGCQQIMEQLQSALKTSEELKSLFNTVATLYCVHQRIDV
 KDTKEALDKIEEIQNKSKQKTQQA AADTQSSSKVSQNYALKHRA YELEFMATTM
 DPVDPNLEPWNHPGSQPTTPGSKYCYKVCYHCPVCFLNKGLGISYGRKKRRQR
 RGT PQSNKDHQNP KQPIPQTQGIS TGPKESKKKVESKTETDPEEFPIPVGEIYK
 RWIFRDYVDRFYKTLRAIFQSSMTKITLWORPLVERYLKDQQLLTVYYGVVPVW
 KRPQVPLRPM TYKAVDL SHFLKEKGGLILKEPVHGVYHPDIVIYQYMDDLTPGP
 GVRYP LACTPYDINQMLRGPGRAFVTIPNPLLGLD

8/11

Fig.7B.

HIVaET DNA

CCCGCCGCCACCATGCCCATCGTGCAGAACGCCAGGGCCAGATGCACCAGGCCCTGTCC
 CCCCGCACCTGAAACGCCTGGGTGAAGGTGATCGAGGAGAAGGCCTTCTCCCCGAGGTG
 ATCCCCATGTTCTCCGCCCTGTCCGAGGGCGCCACCCCCCAGGACCTGAACATGATGCTG
 AACATCGTGGGCGGCCACCAGGCCGCCATGCAGATGCTGAAGGACACCATCAACGAGGAG
 GCCGCCGAGTGGGACCGCCTGCACCCCGTGCACGCCGGCCCCATCCCCCGGCCAGATG
 CGCGAGCCCCGCGGATCCGACATCGCCGGCACCACCTCCACCCTGCAGGAGCAGATCGGC
 TGGATGACCTCCAACCCCCCATCCCCGTGGGCGACATCTACAAGCGTGGATCATCCTG
 GGCCTGAACAAGATCGTGCGCATGTACTCCCCCGTGTCCATCCTGGACATCCGCCAGGGC
 CCAAGGAGCCCTTCCGCGACTACGTGGACCGCTTCTTCAAGACCTGCGCGCCGAGCAG
 GCCACCCAGGAGGTGAAGAACTGGATGACCGAGACCTGCTGGTGCAGAACGCCAACCCC
 GACTGCAAGTCCATCCTGCGCGCCCTGGGCCCCGCGGCCACCCTGGAGGAGATGATGACC
 GCCTGCCAGGGCGTGGGCGGCCCGGCCACAAGGCCCGCGTGTGGGTACCGGCGCCCGC
 GCCTCCGTGCTGTCCGGCGGCAAGCTGGACGCTGGGAGAAGATCCGCCTGCGCCCCGGC
 GGCAAGAAGAAGTACCGCCTGAAGCACCTGGTGTGGGCCTCCCGCGAGCTGGAGCGCTTC
 GCCCTGAACCCCTCCCTGCTGGAGACCGCCGAGGGCTGCCAGCAGATCATGGAGCAGCTG
 CAGTCCGCCCTGAAGACCTCCGAGGAGCTGAAGTCCCTGTTCAACACCGTGGCCACCCTG
 TACTGCGTGCAACAGCGCATCGACGTGAAGGACACCAAGGAGGCCCTGGACAAGATCGAG
 GAGATCCAGAACAAGTCCAAGCAGAAGACCCAGCAGGCCCGCCGACACCCAGTCCTCC
 TCCAAGGTGTCCAGAACTACGCCCTGAAGCACCGCGCCTACGAGCTGGAATTCCCTCCA
 ATTCTGTGCGGGGAGATTTATAAACGGTGGATCATTTTTAGGGATTATGTGATAGGTTT
 TATAAACGCTCAGGGCCATCTTCCAGTCTTCCATGACCAAGATCACCTGTGGCAGCGC
 CCCCTGGTGGAGCGCTACCTGAAGGACCAGCAGCTGCTGACCGTGTACTACGGCGTGCCC
 GTGTGGAAGCGCCCCCAGGTGCCCTGCGCCCCATGACCTACAAGGCCGTGGACCTGTCC
 CACTTCTGAAGGAGAAGGGCGGCCTGATCCTGAAGGAGCCCCGTGCACGGCGTGTACCAC
 CCCGACATCGTGATCTACCAGTACATGGACGACCTGACCCCCGGCCCCGGCGTGCCTAC
 CCCCTGGCCTGCACCCCTACGACATCAACCAGATGCTGCGCGGCCCGGCCGCGCCTTC
 GTGACCATCCCCAACCCCTGCTGGGCCTGGACTGAGCGGCCGCCCTCTCCCTCCCCC
 CCCCTAACGTTACTGGCCGAAGCCGCTTGAATAAGGCCGGTGTGCGTTTGTCTATATGT
 TATTTTCCACCATATTGCCGTCTTTTGGCAATGTGAGGGCCCGGAAACCTGGCCCTGTCT
 TCTTGACGAGCATTCCTAGGGGTCTTTCCCTCTCGCCAAAGGAATGCAAGGTCTGTGA
 ATGTCGTGAAGGAAGCAGTTCCTCTGGAAGCTTCTTGAAGACAAACAACGTCTGTAGCGA
 CCCTTTGCAGGCAGCGGAACCCCCACCTGGCGACAGGTGCCTCTGCGGCCAAAAGCCAC
 GTGTATAAGATACACCTGCAAGGCGGCACAACCCAGTGCCACGTTGTGAGTTGGATAG
 TTGTGGAAGAGTCAAATGGCTCTCCTCAAGCGTATTCAACAAGGGGCTGAAGGATGCC
 AGAAGGTACCCCATTTGTATGGGATCTGATCTGGGGCCTCGGTGCACATGCTTTACATGTG
 TTTAGTCGAGGTTAAAAAACGTCTAGGCCCCCCGAACCACGGGGACGTGGTTTTCCTTT
 GAAAAACACGATGATAATATGGCCACAACCATGGACCCCGTGGACCCCAACCTGGAGCCC
 TGGAACCACCCCGGCTCCAGCCCACCACCCCGGCTCCAAGTGCTACTGCAAGGTGTGC
 TGCTACCACTGCCCCGTGTGCTTCTGAAACAAGGGCCTGGGCATCTCCTACGGCCGCAAG
 AAGCGCCGCCAGCGCCGCGCACCCCCCAGTCCAACAAGGACCACCAGAACCCCATCCCC
 AAGCAGCCCATCCCCAGACCCAGGGCATCTCCACCGGCCCAAGGAGTCCAAGAAGAAG
 GTGGAGTCCAAGACCGAGACCGACCCCGAGTAA

Fig.7A.

HIVaET protein – TAT (HIVa is the same as above)

MATTMDPVDPNLEPWNHPGSQPTTPGSKCYCKVCCYHCPVCFLNKGLGISYGR
 KKRRQRRGTPQSNKDHQNPQKQPIQQTQISTGPKESKKKVESKTETDPE

9/11

Fig.8A.

PPA protein

MPIVQNAQQGMHQALSPRTLNAWVKVIEEKAFSPEVIPMFSALSEGATPQDLNM
MLNIVGGHQAAMQMLKDTINEEAAEWDRHPVHAGPIPPGQMREPRGSDIAGT
TSTLQEQIGWMTSNPPIPVGDYIKRWILGLNKIVRMYSVPSILDIRQGPKEPFRDY
VDRFFKTLRAEQATQEVKNWMTETLLVQANANPDCKSILRALGPGATLEEMMTA
CQGVGGPGHKARVLGTGARASVLSGGKLDWEKIRLRPGGKKKYRLKHLVWA
SRELERFALNPSLLETAEGCQQIMEQLQSALKTSEELKSLFNTVATLYCVHQRIDV
KDTKEALDKIEIQNKSKQKTQAAAADTQSSSKVSQNYALKHRAYELEFGIKVK
QLCKLLRGAKALTDIVTLTEEALELAENREILKDPVHGYYDPSKDLIAEIQKQ
GQDQWTYQIYQEPFKNLKTGKYARKRSAQTNDVKQLAEVVQKVVMESIVIWGK
TPKFRPLPIQKETWETWWMDYWQATWIPEWEFVNTPLVKLWYQLEKDPIAGAE
TFYVDGAANRETKLGKAGYVTDGRGRQKVVSLETETNQKTELHVIHLALQDSGSE
VNIVTDSQYALGIIQAQPDRSDPVDPNLEPWNHPGSQPTTPGSKCYCKVCCYHCP
VCFLNKGGLISYGRKKRRQRRGTPOSNKDHQNPPIKOPIPOTQGISGPKESKKK
VESKTETDPEDAGRSGNSDEELLKAIRIILYQSNPYPKPKGSRQARKNRRRRWR
AGQRQIDSLSERILSTCLGRPAEPVPLQLPPELDCSEDCGTSQTQSQAETGVG
RPQVSVESSAVLGSSTKEGTVRPQVPLRPMTYKAAFDLSFFLKEKGGLDGLIYSK
KRQEILDLWVYHTQGYFPDWQNYTPGPGIRYPLTFGWCFKLVPVDPDEVEEATG
GENNSLLHPICQHGMDDEEKETLRWKFDSSLALKHRARELHPESYKDCPOITLW
QRPLVTKIGGQKTRGGKWSKSSIVGWPEVRERIRQTPTAARERTRQAPTAAKVG
AVSQDLDKHGAVSSNVNHPSCAWLEAQEEEEVGFPPELLDTGADDTVLEDINLPG
KWKPKMIGGIGGLIKVKQYDQILIEICGKKAIGTVLVGPTPVNIIGRNMLTQIGCTL
NFPISPIETVPVKLKPGMDGPKVKQWPLTEEKIKALTEICADMEKEGKISKIGPEN
PYNTPIFAIKKKQSTKWRKLVDFRELNRKTQDFWEVQLGIPHPAGLKKKKS MTV
LDVGDAYFSVPLDESFRKYTAFTIPSTNNETPGVRYQYNVLPQGWKGSPIFQSSM
TKILEPFRSKNPDIVIYQYMDDL YVGS DLEIGQHRTKIEELRAHLLSWGFITPDKK
HQKEPPFLWMGYELHPDKWTVQPIELPEKDSWTVNDIQKLVGKLNWASQIYAC
TPYDINQMLRGPGRAFVTIPNPLLGLD

10/11

Fig.8B.

PPA DNA

CCCCGCCACCATGCCCATCGTGCAGAACGCCAGGGCCAGATGCACCAGGCCCTGTCC
CCCCGCACCCTGAACGCCTGGGTGAAGGTGATCGAGGAGAAGGCCTTCTCCCCCGAGGTG
ATCCCCATGTTCTCCGCCCTGTCCGAGGGCGCCACCCCCAGGACCTGAACATGATGCTG
AACATCGTGGGCGGCCACCAGGCCCATGCGATGCTGAAGGACACCATCAACGAGGAG
GCCGCCGAGTGGGACCGCCTGCACCCCGTGCACGCCGGCCCCATCCCCCGGCCAGATG
CGCGAGCCCCCGCGGATCCGACATCGCCGGCACCACCTCCACCCTGCAGGAGCAGATCGGC
TGGATGACCTCCAACCCCCCATCCCCGTGGGCGACATCTACAAGCGCTGGATCATCCTG
GGCCTGAACAAGATCGTGCATGTACTCCCCGTGTCCATCCTGGACATCCGCCAGGGC
CCCAAGGAGCCCTTCCGCGACTACGTGGACCGCTTCTTCAAGACCCTGCGCGCCGAGCAG
GCCACCCAGGAGGTGAAGAACTGGATGACCGAGACCCTGCTGGTGCAGAACGCCAACCCC
GACTGCAAGTCCATCCTGCGCGCCCTGGGCCCCGGCGCCACCCTGGAGGAGATGATGACC
GCCTGCCAGGGCGTGGGCGGCCCGGCCACAAGGCCCGCGTGTGGGTACCGGCGCCCGC
GCCTCCGTGCTGTCCGGCGGCAAGCTGGACGCTGGGAGAAGATCCGCTGCGCCCCGGC
GGCAAGAAGAAGTACCGCCTGAAGCACCTGGTGTGGGCCTCCGCGAGCTGGAGCGCTT
GCCCTGAACCCCTCCCTGCTGGAGACCGCCGAGGGCTGCCAGCAGATCATGGAGCAGCTG
CAGTCCGCCCTGAAGACCTCCGAGGAGCTGAAGTCCCTGTTCAACACCGTGGCCACCCTG
TACTGCGTGCACCAGCGCATCGACGTGAAGGACACCAAGGAGGCCCTGGACAAGATCGAG
GAGATCCAGAACAAGTCCAAGCAGAAGACCCAGCAGGCCCGCCGCGACACCCAGTCTCTC
TCCAAGGTGTCCAGAACTACGCCCTGAAGCACCGCGCCTACGAGCTGGAATTCCGGCATC
AAGGTGAAGCAGCTGTGCAAGCTGCTGCGCGGCCCAAGGCCCTGACCGACATCGTGACC
CTGACCGAGGAGGCCGAGCTGGAGCTGGCCGAGAACC GCGAGATCCTGAAGGACCCCGTG
CACGGCGTGTACTACGACCCCTCCAAGGACCTGATCGCCGAGATCCAGAAGCAGGGCCAG
GACCAGTGGACCTACCAAATCTACCAGGAGCCCTTCAAGAACCTGAAGACCGGCAAGTAC
GCCCAGAACGCGTCCGCCCAGACCAACGACGTGAAGCAGCTGGCCGAGGTGGTGCAGAAG
GTGGTGAAGAGTCCATCGTGATCTGGGCGAAGACCCCAAGTTCCGCCTGCCCATCCAG
AAGGAGACCTGGGAGACCTGGTGGATGGACTACTGGCAGGCCACCTGGATTCCCGAGTGG
GAGTTCTGTGAACACCCACCCCTGGTGAAGCTGTGGTATCAGCTGGAGAAGGACCCCATC
GCCGGCGCCGAGACCTTCTACGTGGACGGCGCCGCCAACCGCGAGACCAAGCTGGGCAAG
GCCGGCTACGTGACCGACCGGGGCCCGCAGAAGGTGGTGTCCCTGACCGAGACCACCAAC
CAGAAGACCGAGCTGCACGTATCCACCTGGCCCTGCAGGACTCCGGCTCCGAGGTGAAC
ATCGTGACCGACTCCCAGTACGCCCTGGGCATCATCCAGGCCCAGCCCGACAGATCTGAC
CCCGTGGACCCCAACCTGGAGCCCTGGAACCACCCCGGCTCCAGCCACCAACCCCGGC
TCCAAGTGCTACTGCAAGGTGTGCTGTACCACTGCCCCGTGTGCTTCTGAACAAGGGC
CTGGGCATCTCTACGGCCGCAAGAAGCGCCGCGCAGCGCCGCGGCACCCCCAGTCCAAC
AAGGACCAACAGAACCCCATCCCCAAGCAGCCCATCCCCCAGACCCAGGGCATCTCCACC
GGCCCCAAGGAGTCCAAGAAGAAGGTGGAGTCCAAGACCGAGACCCCGAGGACGCC
GGCCGCTCCGGCAACTCCGACGAGGAGCTGCTGAAGGCCATCCGCATCATCAAGATCCTG
TACCAGTCCAACCCCTACCCCAAGCCCAAGGGCTCCCGCCAGGCCCGCAAGAACCGCCGC
CGCCGCTGGCGCGCCGGCCAGCGCCAGATCGACTCCCTGTCCGAGCGCATCCTGTCCACC
TGCCTGGGCCGCCCGCCGAGCCCGTGGCCCTGCAGCTGCCCCCCTGGAGCTGGACTGC
TCCGAGGACTGCGGCACCTCCGGCACCCAGCAGTCCAGGGCGCCGAGACCGGCGTGGGC
CGCCCCCAGGTGTCCGTGGAGTCCCTCCGCCGTGTGGGCTCCGGCACCAAGGAGGGTACC
GTGCGCCCCCAGGTGCCCCGTGCGCCCCATGACCTACAAGGCCGCTTCGACCTGTCTTCT
TTTCTGAAGGAGAAGGGCGGCCTGGACGGCTGATCTACTCCAAGAAGCGCCAGGAGATC
CTGGACCTGTGGGTGTACCACACCCAGGGCTACTTCCCGACTGGCAGAAGTACACCCCC
GGCCCCGGCTCCGCTACCCCTGACCTTCCGCTGGTGTCTTCAAGCTGTGCCCCGTGGAC
CCCGACGAGGTGGAGGAGGCCACCGCGCGGAGAACAACTCCCTGCTGCACCCCATCTGC
CAGCACGGCATGGACGACGAGGAGAAGGAGACCCTGCGCTGGAAGTTCGACTCCTCCCTG
GCCCTGAAGCACCGCGCCCGCAACTCCACCCGAGTCTACAAGGACTGCCCCCAGATC
ACCCTGTGGCAGCGCCCCCTGGTGACCAAGATCGGCGGCCAGAAGACGCGTGGC

11/11

Fig.8B(Cont.)

GGCAAGTGGTCCAAGTCCTCCATCGTGGGCTGGCCCCGAGGTGCGCGAGCGCATCCGCCAG
ACCCCCACCGCCCGCCGCGAGCGCACCCGCCAGGCCCCACCGCCGCCAAGGTGGGCGCC
GTGTCCCAGGACCTGGACAAGCACGGCGCCGTGTCTCCAACGTGAACCACCCCTCCTGC
GCCTGGCTGGAGGCCAGGAAGAGGAAGAGGTGGGCTTCCCCGAGCTCCTGGACACCGGC
GCCGACGACACCGTGCTGGAGGACATCAACCTGCCCGGCAAGTGGAAGCCCCAAGATGATC
GGCGGCATCGGCGGCTTGATCAAGGTGAAGCAGTACGACCAGATCCTGATCGAAATCTGC
GGCAAGAAGGCCATCGGCACCGTGCTGGTGGGCCCCACCCCGTGAACATCATCGGCCGC
AACATGCTGACCCAGATCGGCTGCACCCTGAACTTCCCCATCTCCCCATCGAGACCGTG
CCCGTGAAGCTGAAGCCCGGCATGGACGGCCCCAAGGTGAAGCAGTGGCCCCCTGACCGAG
GAGAAGATCAAGGCCCTGACCGAAATCTGCGCCGACATGGAGAAGGAGGGCAAGATCAGT
AAGATCGGCCCCGAGAACCCCTACAACACCCCATCTTCGCCATCAAGAAGAAGCAGTCC
ACCAAGTGGCGCAAGCTGGTGGACTTCCGCGAGCTGAACAAGCGCACCCAGGACTTCTGG
GAGGTGCAGCTGGGCATCCCCACCCCGCCGCTGAAGAAGAAAAAGTCCGTGACCGTG
CTGGACGTGGGCGACGCCTACTTCTCCGTGCCCCCTGGACGAGTCCTTCCGCAAGTACACC
GCCTTCACCATCCCCCTCCACCAACAACGAGACCCCCGGCGTGCGCTACCAGTACAACGTG
CTGCCCCAGGGCTGGAAGGGATCCCCCATCTTCCAGTCCTCCATGACCAAGATCCTGGAG
CCCTTCCGCTCCAAGAACCCGACATCGTGATCTACCAGTACATGGACGACCTGTACGTG
GGCTCCGACCTGGAGATCGGCCAGCACCGCACCAAGATCGAGGAGCTGCGCGCCCACCTG
CTGTCTGGGGCTTCATACCCCCGACAAGAAGCACCAGAAGGAGCCCCCTTCTGTGG
ATGGGCTACGAGCTGCACCCGACAAGTGGACCGTGACGCCATCGAGCTGCCCGAGAAG
GACTCCTGGACCGTGAACGACATCCAGAAGCTGGTGGGCAAGCTGAACTGGGCCTCCCAA
ATCTACGCCTGCACCCCTACGACATCAACCAGATGCTGCGCGGCCCCGGCCGCGCCTTC
GTGACCATCCCCAACCCCTGCTGGGCTGGACTA

SEQUENCE LISTING

<110> Medical Research Council
 International Aids Vaccine Initiative
 University of Nairobi

<120> Improvements in or Relating to Immune Responses to HIV

<130> MJL/C1248'1/M

<140>

<141>

<160> 33

<170> PatentIn Ver. 2.1

<210> 1

<211> 11

<212> PRT

<213> Simian immunodeficiency virus

<400> 1

Ala	Cys	Thr	Pro	Tyr	Asp	Ile	Asn	Gln	Met	Leu
1					5				10	

<210> 2

<211> 10

<212> PRT

<213> Human immunodeficiency virus

<400> 2

Arg	Gly	Pro	Gly	Arg	Ala	Phe	Val	Thr	Ile
1				5					10

<210> 3

<211> 9

<212> PRT

<213> Human immunodeficiency virus

<400> 3

Ala Leu Lys His Arg Ala Tyr Glu Leu
1 5

<210> 4

<211> 9

<212> PRT

<213> Human immunodeficiency virus

<400> 4

Pro Pro Ile Pro Val Gly Glu Ile Tyr
1 5

<210> 5

<211> 9

<212> PRT

<213> Human immunodeficiency virus

<400> 5

Gly Glu Ile Tyr Lys Arg Trp Ile Ile
1 5

<210> 6

<211> 10

<212> PRT

<213> Human immunodeficiency virus

<400> 6

Lys Arg Trp Ile Ile Leu Gly Leu Asn Lys
1 5 10

<210> 7

<211> 10

<212> PRT

<213> Human immunodeficiency virus

<400> 7

Phe Arg Asp Tyr Val Asp Arg Phe Tyr Lys
1 5 10

<210> 8
<211> 11
<212> PRT
<213> Human immunodeficiency virus

<400> 8
Arg Asp Tyr Val Asp Arg Phe Tyr Lys Thr Leu
1 5 10

<210> 9
<211> 9
<212> PRT
<213> Human immunodeficiency virus

<400> 9
Asp Arg Phe Tyr Lys Thr Leu Arg Ala
1 5

<210> 10
<211> 9
<212> PRT
<213> Human immunodeficiency virus

<400> 10
Ala Ile Phe Gln Ser Ser Met Thr Lys
1 5

<210> 11
<211> 9
<212> PRT
<213> Human immunodeficiency virus

<400> 11
Ile Thr Leu Trp Gln Arg Pro Leu Val
1 5

<210> 12
<211> 9
<212> PRT
<213> Human immunodeficiency virus

<400> 12

Glu Arg Tyr Leu Lys Asp Gln Gln Leu
1 5

<210> 13

<211> 8

<212> PRT

<213> Human immunodeficiency virus

<400> 13

Tyr Leu Lys Asp Gln Gln Leu Leu
1 5

<210> 14

<211> 10

<212> PRT

<213> Human immunodeficiency virus

<400> 14

Thr Val Tyr Tyr Gly Val Pro Val Trp Lys
1 5 10

<210> 15

<211> 11

<212> PRT

<213> Human immunodeficiency virus

<400> 15

Arg Pro Gln Val Pro Leu Arg Pro Met Thr Tyr
1 5 10

<210> 16

<211> 10

<212> PRT

<213> Human immunodeficiency virus

<400> 16

Gln Val Pro Leu Arg Pro Met Thr Tyr Lys
1 5 10

<210> 17
<211> 8
<212> PRT
<213> Human immunodeficiency virus

<400> 17
Val Pro Leu Arg Pro Met Thr Tyr
1 5

<210> 18
<211> 9
<212> PRT
<213> Human immunodeficiency virus

<400> 18
Ala Val Asp Leu Ser His Phe Leu Lys
1 5

<210> 19
<211> 9
<212> PRT
<213> Human immunodeficiency virus

<400> 19
Asp Leu Ser His Phe Leu Lys Glu Lys
1 5

<210> 20
<211> 8
<212> PRT
<213> Human immunodeficiency virus

<400> 20
Phe Leu Lys Glu Lys Gly Gly Leu
1 5

<210> 21
<211> 9
<212> PRT
<213> Human immunodeficiency virus

<400> 21

Ile Leu Lys Glu Pro Val His Gly Val

1

5

<210> 22

<211> 10

<212> PRT

<213> Human immunodeficiency virus

<400> 22

Ile Leu Lys Glu Pro Val His Gly Val Tyr

1

5

10

<210> 23

<211> 9

<212> PRT

<213> Human immunodeficiency virus

<400> 23

His Pro Asp Ile Val Ile Tyr Gln Tyr

1

5

<210> 24

<211> 9

<212> PRT

<213> Human immunodeficiency virus

<400> 24

Val Ile Tyr Gln Tyr Met Asp Asp Leu

1

5

<210> 25

<211> 10

<212> PRT

<213> Human immunodeficiency virus

<400> 25

Thr Pro Gly Pro Gly Val Arg Tyr Pro Leu

1

5

10

<210> 26
 <211> 527
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Chimeric
 polypeptide

<400> 26
 Met Pro Ile Val Gln Asn Ala Gln Gly Gln Met His Gln Ala Leu Ser
 1 5 10 15
 Pro Arg Thr Leu Asn Ala Trp Val Lys Val Ile Glu Glu Lys Ala Phe
 20 25 30
 Ser Pro Glu Val Ile Pro Met Phe Ser Ala Leu Ser Glu Gly Ala Thr
 35 40 45
 Pro Gln Asp Leu Asn Met Met Leu Asn Ile Val Gly Gly His Gln Ala
 50 55 60
 Ala Met Gln Met Leu Lys Asp Thr Ile Asn Glu Glu Ala Ala Glu Trp
 65 70 75 80
 Asp Arg Leu His Pro Val His Ala Gly Pro Ile Pro Pro Gly Gln Met
 85 90 95
 Arg Glu Pro Arg Gly Ser Asp Ile Ala Gly Thr Thr Ser Thr Leu Gln
 100 105 110
 Glu Gln Ile Gly Trp Met Thr Ser Asn Pro Pro Ile Pro Val Gly Asp
 115 120 125
 Ile Tyr Lys Arg Trp Ile Ile Leu Gly Leu Asn Lys Ile Val Arg Met
 130 135 140
 Tyr Ser Pro Val Ser Ile Leu Asp Ile Arg Gln Gly Pro Lys Glu Pro
 145 150 155 160
 Phe Arg Asp Tyr Val Asp Arg Phe Phe Lys Thr Leu Arg Ala Glu Gln
 165 170 175
 Ala Thr Gln Glu Val Lys Asn Trp Met Thr Glu Thr Leu Leu Val Gln
 180 185 190

Asn Ala Asn Pro Asp Cys Lys Ser Ile Leu Arg Ala Leu Gly Pro Gly
 195 200 205
 Ala Thr Leu Glu Glu Met Met Thr Ala Cys Gln Gly Val Gly Gly Pro
 210 215 220
 Gly His Lys Ala Arg Val Leu Gly Thr Gly Ala Arg Ala Ser Val Leu
 225 230 235 240
 Ser Gly Gly Lys Leu Asp Ala Trp Glu Lys Ile Arg Leu Arg Pro Gly
 245 250 255
 Gly Lys Lys Lys Tyr Arg Leu Lys His Leu Val Trp Ala Ser Arg Glu
 260 265 270
 Leu Glu Arg Phe Ala Leu Asn Pro Ser Leu Leu Glu Thr Ala Glu Gly
 275 280 285
 Cys Gln Gln Ile Met Glu Gln Leu Gln Ser Ala Leu Lys Thr Ser Glu
 290 295 300
 Glu Leu Lys Ser Leu Phe Asn Thr Val Ala Thr Leu Tyr Cys Val His
 305 310 315 320
 Gln Arg Ile Asp Val Lys Asp Thr Lys Glu Ala Leu Asp Lys Ile Glu
 325 330 335
 Glu Ile Gln Asn Lys Ser Lys Gln Lys Thr Gln Gln Ala Ala Ala Asp
 340 345 350
 Thr Gln Ser Ser Ser Lys Val Ser Gln Asn Tyr Ala Leu Lys His Arg
 355 360 365
 Ala Tyr Glu Leu Glu Phe Pro Pro Ile Pro Val Gly Glu Ile Tyr Lys
 370 375 380
 Arg Trp Ile Ile Phe Arg Asp Tyr Val Asp Arg Phe Tyr Lys Thr Leu
 385 390 395 400
 Arg Ala Ile Phe Gln Ser Ser Met Thr Lys Ile Thr Leu Trp Gln Arg
 405 410 415
 Pro Leu Val Glu Arg Tyr Leu Lys Asp Gln Gln Leu Leu Thr Val Tyr
 420 425 430

Tyr Gly Val Pro Val Trp Lys Arg Pro Gln Val Pro Leu Arg Pro Met
 435 440 445

Thr Tyr Lys Ala Val Asp Leu Ser His Phe Leu Lys Glu Lys Gly Gly
 450 455 460

Leu Ile Leu Lys Glu Pro Val His Gly Val Tyr His Pro Asp Ile Val
 465 470 475 480

Ile Tyr Gln Tyr Met Asp Asp Leu Thr Pro Gly Pro Gly Val Arg Tyr
 485 490 495

Pro Leu Ala Cys Thr Pro Tyr Asp Ile Asn Gln Met Leu Arg Gly Pro
 500 505 510

Gly Arg Ala Phe Val Thr Ile Pro Asn Pro Leu Leu Gly Leu Asp
 515 520 525

<210> 27

<211> 1608

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Chimeric polynucleotide

<400> 27

```

aagcttcccc cgcaccacat gccatcgtg cagaacgccc agggccagat gcaccaggcc 60
ctgtcccccc gcaccctgaa cgcctgggtg aaggtgatcg aggagaaggc cttctcccc 120
gaggtgatcc ccatgttctc cgccctgtcc gagggcgcca cccccagga cctgaacatg 180
atgctgaaca tcgtgggcgg ccaccaggcc gccatgcaga tgctgaagga caccatcaac 240
gaggaggccg ccgagtggga ccgcctgcac ccctgtcacg ccggccccc ccccccgcc 300
cagatgcgcg agccccgcgg atccgacatc gccggcacca cctccaccct gcaggagcag 360
atcggtgga tgacctcaa ccccccatc ccctgtggcg acatctacaa gcgctggatc 420
atcctgggcc tgaacaagat cgtacgcag tactccccg tgtccatcct ggacatccgc 480
cagggcccca aggagccctt ccgcgactac gtggaccgct tcttcaagac cctgcgcgcc 540
gagcaggcca cccaggaggt gaagaactgg atgaccgaga ccctgctggt gcagaacgcc 600
aaccgcgact gcaagtccat cctgcgcgcc ctgggccccg gcgccaccct ggaggagatg 660
atgaccgcct gccagggcgt gggcgggccc ggccacaagg ccgcgtgct gggtagccgc 720
gccgcgcct ccgtgctgtc cggcggaag ctggacgcct gggagaagat ccgcctgcgc 780
cccgcgcca agaagaagta ccgcctgaag cacctggtgt gggcctccc cgagctggag 840
cgcttcgccc tgaaccctc cctgctggag accgccgagg gctgccagca gatcatggag 900
cagctgcagt ccgcctgaa gacctccgag gagctgaagt ccctgttcaa caccgtggcc 960

```

```

accctgtact gcgtgcacca gcgcatcgac gtgaaggaca ccaaggaggc cctggacaag 1020
atcgaggaga tccagaacaa gtccaagcag aagacccagc aggccgccgc cgacaccag 1080
tcctcctcca aggtgtcca gaactacgcc ctgaagcacc gcgcctacga gctggaattc 1140
cctccaattc ctgtcgggga gatttataaa cggtggatca ttttaggga ttatgtcgat 1200
aggttttata aaacgctcag ggccatcttc cagtcctcca tgaccaagat caccctgtgg 1260
cagcgccccc tgggtggagcg ctacctgaag gaccagcagc tgctgaccgt gtactacggc 1320
gtgcccgtgt ggaagcgccc ccaggtgccc ctgcgcccca tgacctacaa ggccgtggac 1380
ctgtcccact tcctgaagga gaaggcggc ctgacccga aggagcccgt gcacggcgtg 1440
taccaccccg acatcgtgat ctaccagtac atggacgacc tgacccccgg ccccggcgtg 1500
cgctaccccc tggcctgcac cccctacgac atcaaccaga tgctgcgcgg ccccgccgcg 1560
gccttcgtga ccatcccaaa cccctgctg ggcctggact gatctaga 1608

```

<210> 28

<211> 633

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Chimeric
polypeptide

<400> 28

```

Met Pro Ile Val Gln Asn Ala Gln Gly Gln Met His Gln Ala Leu Ser
  1             5             10             15

```

```

Pro Arg Thr Leu Asn Ala Trp Val Lys Val Ile Glu Glu Lys Ala Phe
          20             25             30

```

```

Ser Pro Glu Val Ile Pro Met Phe Ser Ala Leu Ser Glu Gly Ala Thr
          35             40             45

```

```

Pro Gln Asp Leu Asn Met Met Leu Asn Ile Val Gly Gly His Gln Ala
          50             55             60

```

```

Ala Met Gln Met Leu Lys Asp Thr Ile Asn Glu Glu Ala Ala Glu Trp
          65             70             75             80

```

```

Asp Arg Leu His Pro Val His Ala Gly Pro Ile Pro Pro Gly Gln Met
          85             90             95

```

```

Arg Glu Pro Arg Gly Ser Asp Ile Ala Gly Thr Thr Ser Thr Leu Gln
          100            105            110

```

Glu Gln Ile Gly Trp Met Thr Ser Asn Pro Pro Ile Pro Val Gly Asp
 115 120 125
 Ile Tyr Lys Arg Trp Ile Ile Leu Gly Leu Asn Lys Ile Val Arg Met
 130 135 140
 Tyr Ser Pro Val Ser Ile Leu Asp Ile Arg Gln Gly Pro Lys Glu Pro
 145 150 155 160
 Phe Arg Asp Tyr Val Asp Arg Phe Phe Lys Thr Leu Arg Ala Glu Gln
 165 170 175
 Ala Thr Gln Glu Val Lys Asn Trp Met Thr Glu Thr Leu Leu Val Gln
 180 185 190
 Asn Ala Asn Pro Asp Cys Lys Ser Ile Leu Arg Ala Leu Gly Pro Gly
 195 200 205
 Ala Thr Leu Glu Glu Met Met Thr Ala Cys Gln Gly Val Gly Gly Pro
 210 215 220
 Gly His Lys Ala Arg Val Leu Gly Thr Gly Ala Arg Ala Ser Val Leu
 225 230 235 240
 Ser Gly Gly Lys Leu Asp Ala Trp Glu Lys Ile Arg Leu Arg Pro Gly
 245 250 255
 Gly Lys Lys Lys Tyr Arg Leu Lys His Leu Val Trp Ala Ser Arg Glu
 260 265 270
 Leu Glu Arg Phe Ala Leu Asn Pro Ser Leu Leu Glu Thr Ala Glu Gly
 275 280 285
 Cys Gln Gln Ile Met Glu Gln Leu Gln Ser Ala Leu Lys Thr Ser Glu
 290 295 300
 Glu Leu Lys Ser Leu Phe Asn Thr Val Ala Thr Leu Tyr Cys Val His
 305 310 315 320
 Gln Arg Ile Asp Val Lys Asp Thr Lys Glu Ala Leu Asp Lys Ile Glu
 325 330 335
 Glu Ile Gln Asn Lys Ser Lys Gln Lys Thr Gln Gln Ala Ala Ala Asp
 340 345 350

Thr Gln Ser Ser Ser Lys Val Ser Gln Asn Tyr Ala Leu Lys His Arg
 355 360 365
 Ala Tyr Glu Leu Glu Phe Met Ala Thr Thr Met Asp Pro Val Asp Pro
 370 375 380
 Asn Leu Glu Pro Trp Asn His Pro Gly Ser Gln Pro Thr Thr Pro Gly
 385 390 395 400
 Ser Lys Cys Tyr Cys Lys Val Cys Cys Tyr His Cys Pro Val Cys Phe
 405 410 415
 Leu Asn Lys Gly Leu Gly Ile Ser Tyr Gly Arg Lys Lys Arg Arg Gln
 420 425 430
 Arg Arg Gly Thr Pro Gln Ser Asn Lys Asp His Gln Asn Pro Ile Pro
 435 440 445
 Lys Gln Pro Ile Pro Gln Thr Gln Gly Ile Ser Thr Gly Pro Lys Glu
 450 455 460
 Ser Lys Lys Lys Val Glu Ser Lys Thr Glu Thr Asp Pro Glu Glu Phe
 465 470 475 480
 Pro Pro Ile Pro Val Gly Glu Ile Tyr Lys Arg Trp Ile Ile Phe Arg
 485 490 495
 Asp Tyr Val Asp Arg Phe Tyr Lys Thr Leu Arg Ala Ile Phe Gln Ser
 500 505 510
 Ser Met Thr Lys Ile Thr Leu Trp Gln Arg Pro Leu Val Glu Arg Tyr
 515 520 525
 Leu Lys Asp Gln Gln Leu Leu Thr Val Tyr Tyr Gly Val Pro Val Trp
 530 535 540
 Lys Arg Pro Gln Val Pro Leu Arg Pro Met Thr Tyr Lys Ala Val Asp
 545 550 555 560
 Leu Ser His Phe Leu Lys Glu Lys Gly Gly Leu Ile Leu Lys Glu Pro
 565 570 575
 Val His Gly Val Tyr His Pro Asp Ile Val Ile Tyr Gln Tyr Met Asp
 580 585 590

Asp Leu Thr Pro Gly Pro Gly Val Arg Tyr Pro Leu Ala Cys Thr Pro
 595 600 605

Tyr Asp Ile Asn Gln Met Leu Arg Gly Pro Gly Arg Ala Phe Val Thr
 610 615 620

Ile Pro Asn Pro Leu Leu Gly Leu Asp
 625 630

<210> 29

<211> 1914

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Chimeric polynucleotide

<400> 29

```

ccccgcccca ccatgcccat cgtgcagaac gcccagggcc agatgcacca ggccctgtcc 60
ccccgcaccc tgaacgcctg ggtgaagggtg atcgaggaga aggccttctc ccccagagggtg 120
atcccatgtg tctccgcctt gtccgagggc gccaccccc aggacctgaa catgatgctg 180
aacatcgtgg gcggccacca ggccgccatg cagatgctga aggacacat caacgaggag 240
gccgccgagt gggaccgcct gcaccccggt cagcgcggcc ccaccccc cgccagatg 300
cgcgagcccc gcggatccga catcgccggc accacctcca ccctgcagga gcagatcggc 360
tggtatgacct ccaaccccc catcccggtg ggcgacatct acaagcgctg gatcatcctg 420
ggcctgaaca agatcgtgct catgtactcc cccgtgtcca tcctggacat ccgccagggc 480
cccaaggagc ccttcgcga ctacgtggac cgcttcttca agaccctgcg cgccgagcag 540
gccaccagg aggtgaagaa ctggatgacc gagaccctgc tgggtgcagaa cgccaacccc 600
gactgcaagt ccacctgcg cgccctgggc cccggcgcca ccctggagga gatgatgacc 660
gcctgccagg gcgtgggcgg cccggccac aaggcccgcg tgcgggtac cggcgcccgc 720
gcctccgtgc tgcgcggcg caagctggac gcctgggaga agatccgcct gcgccccggc 780
ggcaagaaga agtaccgcct gaagcacctg gtgtgggcct cccgcgagct ggagcgcttc 840
gccctgaacc cctccctgct ggagaccgcc gagggctgcc agcagatcat ggagcagctg 900
cagtccgccc tgaagacctc cgaggagctg aagtcctgt tcaacaccgt ggccaccctg 960
tactgcgtgc accagcgcac cgacgtgaag gacaccaagg aggccttga caagatcgag 1020
gagatccaga acaagtccaa gcagaagacc cagcaggccg ccgccgacac ccagtcctcc 1080
tccaaggtgt cccagaacta cgccctgaag caccgcgcct acgagctgga attcatggcc 1140
acaaccatgg acccctgga ccccaacctg gagccctgga accaccccg ctccagccc 1200
accaccccc gctccaagt ctactgcaag gtgtgctgct accactgccc cgtgtgcttc 1260
ctgaacaagg gcctgggcat ctctacggc cgcaagaagc gccgccagcg ccgcggcacc 1320
ccccagtcca acaaggacca ccagaacccc atccccaagc agcccatccc ccagaccag 1380
ggcatctcca ccggtcccaa ggagtccaag aagaagggtg agtccaagac cgagaccgac 1440
cccgaggaaat tccctccaat tcctgtcggg gagatttata aacggtggat cattttttag 1500

```

```

gattatgtcg ataggtttta taaaacgctc agggccatct tccagtcctc catgaccaag 1560
atcacctgt ggcagcgccc cctggtggag cgctacctga aggaccagca gctgctgacc 1620
gtgtactacg gcgtgcccgt gtggaagcgc ccccaggtgc ccctgcgcc catgacctac 1680
aaggccgtgg acctgtccca cticctgaag gagaagggcg gcctgaccc gaaggagccc 1740
gtgcacggcg tgtaccaccc cgacatcgtg atctaccagt acatggacga cctgaccccc 1800
ggccccggcg tgcgctaccc cctggcctgc accccctacg acatcaacca gatgctgcgc 1860
ggccccggcg gcgccttcgt gaccatcccc aaccctcgc tgggcctgga ctga 1914

```

<210> 30

<211> 104

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Chimeric
polypeptide

<400> 30

```

Met Ala Thr Thr Met Asp Pro Val Asp Pro Asn Leu Glu Pro Trp Asn
  1             5             10             15

```

```

His Pro Gly Ser Gln Pro Thr Thr Pro Gly Ser Lys Cys Tyr Cys Lys
      20             25             30

```

```

Val Cys Cys Tyr His Cys Pro Val Cys Phe Leu Asn Lys Gly Leu Gly
      35             40             45

```

```

Ile Ser Tyr Gly Arg Lys Lys Arg Arg Gln Arg Arg Gly Thr Pro Gln
      50             55             60

```

```

Ser Asn Lys Asp His Gln Asn Pro Ile Pro Lys Gln Pro Ile Pro Gln
      65             70             75             80

```

```

Thr Gln Gly Ile Ser Thr Gly Pro Lys Glu Ser Lys Lys Lys Val Glu
      85             90             95

```

```

Ser Lys Thr Glu Thr Asp Pro Glu
      100

```

<210> 31

<211> 2493

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Chimeric polynucleotide

<400> 31

```

ccccgcccca ccatgcccat cgtgcagaac gcccagggcc agatgcacca ggccctgtcc 60
ccccgcaccc tgaacgcctg ggtgaagggtg atcgaggaga aggccttctc ccccaggtg 120
atccccatgt tctccgccct gtccgagggc gccaccccc aggacctgaa catgatgtg 180
aacatcgtgg gcggccacca ggccgccatg cagatgctga aggacaccat caacgaggag 240
gccgccgagt gggaccgcct gcaccccggtg cagcgggcc ccatcccccc cgccagatg 300
cgcgagcccc gcggatccga catcgccggc accacctcca ccctgcagga gcagatcggc 360
tggatgacct ccaaccccc catccccgtg ggcgacatct acaagcgctg gatcatcctg 420
ggcctgaaca agatcgtgctg catgtactcc cccgtgtcca tcctggacat ccgccagggc 480
cccaaggagc ccttcgcga ctacgtggac cgcttcttca agaccctgcg cgccgagcag 540
gccaccagg aggtgaagaa ctggatgacc gagaccctgc tgggtgcagaa cgccaacccc 600
gactgcaagt ccatcctgcg cgccctgggc cccggcgcca ccctggagga gatgatgacc 660
gcctgccagg gcgtgggcgg ccccgccac aaggcccgcg tgctgggtac cggcgccgcg 720
gcctccgtgc tgtccggcgg caagctggac gcctgggaga agatccgcct gcgccccggc 780
ggcaagaaga agtaccgcct gaagcacctg gtgtgggcct cccgcgagct ggagcgcttc 840
gccctgaacc cctcctgct ggagaccgcc gagggtgcc agcagatcat ggagcagctg 900
cagtcggccc tgaagacctc cgaggagctg aagtcctgt tcaacaccgt ggccaccctg 960
tactgctgct accagcgcct cgacgtgaag gacaccaagg aggccttga caagatcgag 1020
gagatccaga acaagtccaa gcagaagacc cagcaggccg ccgccgacac ccagtcctcc 1080
tccaagggtg cccagaacta cgccctgaag caccgcgcct acgagctgga attccctcca 1140
attcctgtcg gggagattta taaacggtgg atcattttta gggattatgt cgataggttt 1200
tataaaacgc tcagggccat ctccagtc tccatgacca agatcacct gtggcagcgc 1260
cccctggtgg agcgtacct gaaggaccag cagctgctga ccgtgtacta cggcgtgccc 1320
gtgtggaagc gccccaggt gcccctgcgc cccatgacct acaaggccgt ggacctgtcc 1380
cacttcctga aggagaagg cgccctgac ctgaaggagc ccgtgcacgg cgtgtaccac 1440
cccgacatcg tgatctacca gtacatggac gacctgacct ccggccccgg cgtgcgctac 1500
cccctggcct gcaccccta cgacatcaac cagatgctgc gcggccccgg ccgcgccttc 1560
gtgacctcc ccaacccct gctgggcctg gactgagcgg ccgccccctt ccctcccccc 1620
cccctaactg tactggccga agccgcttgg aataaggccg gtgtgcgttt gtctatatgt 1680
tattttccac catattgccg tcttttgca atgtgagggc ccgaaaacct ggccctgtct 1740
tcttgacgag cattcctagg ggtctttccc ctctcgccaa aggaatgcaa ggtctgttga 1800
atgtcgtgaa ggaagcagtt cctctggaag cttcttgaag acaaacaacg tctgtagcga 1860
ccctttgcag gcagcggaac cccccacctg gcgacaggtg cctctgcggc caaaagccac 1920
gtgtataaga tacacctgca aaggcggcac aacccagtg ccacgttggt agttggatag 1980
ttgtggaag agtcaaattg ctctcctcaa gcgtattcaa caaggggctg aaggatgccc 2040
agaaggtacc ccattgtatg ggatctgatc tggggcctcg gtgcacatgc ttacatgtg 2100
tttagtcgag gttaaaaaaa cgtctaggcc ccccgaacca cggggacgtg gttttccttt 2160
gaaaaacacg atgataatat ggccacaacc atggaccccg tggaccccaa cctggagccc 2220
tgaaccacc ccggtccca gcccaccacc cccggctcca agtgctactg caagggtgtg 2280
tgctaccact gccccgtgtg cttcctgaac aagggcctgg gcatctccta cgccgcaag 2340
aagcgccgcc agcgccgcgg cccccccag tccaacaagg accaccagaa cccatcccc 2400

```


aagcagccca tccccagac ccagggcatc tccaccggcc ccaaggagtc caagaagaag 2460
 gtggagtcca agaccgagac cgaccccgag taa 2493

<210> 32

<211> 1445

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Chimeric
 polypeptide

<400> 32

Met Pro Ile Val Gln Asn Ala Gln Gly Gln Met His Gln Ala Leu Ser
 1 5 10 15

Pro Arg Thr Leu Asn Ala Trp Val Lys Val Ile Glu Glu Lys Ala Phe
 20 25 30

Ser Pro Glu Val Ile Pro Met Phe Ser Ala Leu Ser Glu Gly Ala Thr
 35 40 45

Pro Gln Asp Leu Asn Met Met Leu Asn Ile Val Gly Gly His Gln Ala
 50 55 60

Ala Met Gln Met Leu Lys Asp Thr Ile Asn Glu Glu Ala Ala Glu Trp
 65 70 75 80

Asp Arg Leu His Pro Val His Ala Gly Pro Ile Pro Pro Gly Gln Met
 85 90 95

Arg Glu Pro Arg Gly Ser Asp Ile Ala Gly Thr Thr Ser Thr Leu Gln
 100 105 110

Glu Gln Ile Gly Trp Met Thr Ser Asn Pro Pro Ile Pro Val Gly Asp
 115 120 125

Ile Tyr Lys Arg Trp Ile Ile Leu Gly Leu Asn Lys Ile Val Arg Met
 130 135 140

Tyr Ser Pro Val Ser Ile Leu Asp Ile Arg Gln Gly Pro Lys Glu Pro
 145 150 155 160

Phe Arg Asp Tyr Val Asp Arg Phe Phe Lys Thr Leu Arg Ala Glu Gln
 165 170 175

Ala Thr Gln Glu Val Lys Asn Trp Met Thr Glu Thr Leu Leu Val Gln
 180 185 190
 Asn Ala Asn Pro Asp Cys Lys Ser Ile Leu Arg Ala Leu Gly Pro Gly
 195 200 205
 Ala Thr Leu Glu Glu Met Met Thr Ala Cys Gln Gly Val Gly Gly Pro
 210 215 220
 Gly His Lys Ala Arg Val Leu Gly Thr Gly Ala Arg Ala Ser Val Leu
 225 230 235 240
 Ser Gly Gly Lys Leu Asp Ala Trp Glu Lys Ile Arg Leu Arg Pro Gly
 245 250 255
 Gly Lys Lys Lys Tyr Arg Leu Lys His Leu Val Trp Ala Ser Arg Glu
 260 265 270
 Leu Glu Arg Phe Ala Leu Asn Pro Ser Leu Leu Glu Thr Ala Glu Gly
 275 280 285
 Cys Gln Gln Ile Met Glu Gln Leu Gln Ser Ala Leu Lys Thr Ser Glu
 290 295 300
 Glu Leu Lys Ser Leu Phe Asn Thr Val Ala Thr Leu Tyr Cys Val His
 305 310 315 320
 Gln Arg Ile Asp Val Lys Asp Thr Lys Glu Ala Leu Asp Lys Ile Glu
 325 330 335
 Glu Ile Gln Asn Lys Ser Lys Gln Lys Thr Gln Gln Ala Ala Ala Asp
 340 345 350
 Thr Gln Ser Ser Ser Lys Val Ser Gln Asn Tyr Ala Leu Lys His Arg
 355 360 365
 Ala Tyr Glu Leu Glu Phe Gly Ile Lys Val Lys Gln Leu Cys Lys Leu
 370 375 380
 Leu Arg Gly Ala Lys Ala Leu Thr Asp Ile Val Thr Leu Thr Glu Glu
 385 390 395 400
 Ala Glu Leu Glu Leu Ala Glu Asn Arg Glu Ile Leu Lys Asp Pro Val
 405 410 415

His Gly Val Tyr Tyr Asp Pro Ser Lys Asp Leu Ile Ala Glu Ile Gln
 420 425 430

Lys Gln Gly Gln Asp Gln Trp Thr Tyr Gln Ile Tyr Gln Glu Pro Phe
 435 440 445

Lys Asn Leu Lys Thr Gly Lys Tyr Ala Arg Lys Arg Ser Ala Gln Thr
 450 455 460

Asn Asp Val Lys Gln Leu Ala Glu Val Val Gln Lys Val Val Met Glu
 465 470 475 480

Ser Ile Val Ile Trp Gly Lys Thr Pro Lys Phe Arg Leu Pro Ile Gln
 485 490 495

Lys Glu Thr Trp Glu Thr Trp Trp Met Asp Tyr Trp Gln Ala Thr Trp
 500 505 510

Ile Pro Glu Trp Glu Phe Val Asn Thr Pro Pro Leu Val Lys Leu Trp
 515 520 525

Tyr Gln Leu Glu Lys Asp Pro Ile Ala Gly Ala Glu Thr Phe Tyr Val
 530 535 540

Asp Gly Ala Ala Asn Arg Glu Thr Lys Leu Gly Lys Ala Gly Tyr Val
 545 550 555 560

Thr Asp Arg Gly Arg Gln Lys Val Val Ser Leu Thr Glu Thr Thr Asn
 565 570 575

Gln Lys Thr Glu Leu His Val Ile His Leu Ala Leu Gln Asp Ser Gly
 580 585 590

Ser Glu Val Asn Ile Val Thr Asp Ser Gln Tyr Ala Leu Gly Ile Ile
 595 600 605

Gln Ala Gln Pro Asp Arg Ser Asp Pro Val Asp Pro Asn Leu Glu Pro
 610 615 620

Trp Asn His Pro Gly Ser Gln Pro Thr Thr Pro Gly Ser Lys Cys Tyr
 625 630 635 640

Cys Lys Val Cys Cys Tyr His Cys Pro Val Cys Phe Leu Asn Lys Gly
 645 650 655

Leu Gly Ile Ser Tyr Gly Arg Lys Lys Arg Arg Gln Arg Arg Gly Thr
 660 665 670
 Pro Gln Ser Asn Lys Asp His Gln Asn Pro Ile Pro Lys Gln Pro Ile
 675 680 685
 Pro Gln Thr Gln Gly Ile Ser Thr Gly Pro Lys Glu Ser Lys Lys Lys
 690 695 700
 Val Glu Ser Lys Thr Glu Thr Asp Pro Glu Asp Ala Gly Arg Ser Gly
 705 710 715 720
 Asn Ser Asp Glu Glu Leu Leu Lys Ala Ile Arg Ile Ile Lys Ile Leu
 725 730 735
 Tyr Gln Ser Asn Pro Tyr Pro Lys Pro Lys Gly Ser Arg Gln Ala Arg
 740 745 750
 Lys Asn Arg Arg Arg Arg Trp Arg Ala Gly Gln Arg Gln Ile Asp Ser
 755 760 765
 Leu Ser Glu Arg Ile Leu Ser Thr Cys Leu Gly Arg Pro Ala Glu Pro
 770 775 780
 Val Pro Leu Gln Leu Pro Pro Leu Glu Leu Asp Cys Ser Glu Asp Cys
 785 790 795 800
 Gly Thr Ser Gly Thr Gln Gln Ser Gln Gly Ala Glu Thr Gly Val Gly
 805 810 815
 Arg Pro Gln Val Ser Val Glu Ser Ser Ala Val Leu Gly Ser Gly Thr
 820 825 830
 Lys Glu Gly Thr Val Arg Pro Gln Val Pro Leu Arg Pro Met Thr Tyr
 835 840 845
 Lys Ala Ala Phe Asp Leu Ser Phe Phe Leu Lys Glu Lys Gly Gly Leu
 850 855 860
 Asp Gly Leu Ile Tyr Ser Lys Lys Arg Gln Glu Ile Leu Asp Leu Trp
 865 870 875 880
 Val Tyr His Thr Gln Gly Tyr Phe Pro Asp Trp Gln Asn Tyr Thr Pro
 885 890 895

Gly Pro Gly Ile Arg Tyr Pro Leu Thr Phe Gly Trp Cys Phe Lys Leu
 900 905 910

Val Pro Val Asp Pro Asp Glu Val Glu Glu Ala Thr Gly Gly Glu Asn
 915 920 925

Asn Ser Leu Leu His Pro Ile Cys Gln His Gly Met Asp Asp Glu Glu
 930 935 940

Lys Glu Thr Leu Arg Trp Lys Phe Asp Ser Ser Leu Ala Leu Lys His
 945 950 955 960

Arg Ala Arg Glu Leu His Pro Glu Ser Tyr Lys Asp Cys Pro Gln Ile
 965 970 975

Thr Leu Trp Gln Arg Pro Leu Val Thr Lys Ile Gly Gly Gln Lys Thr
 980 985 990

Arg Gly Gly Lys Trp Ser Lys Ser Ser Ile Val Gly Trp Pro Glu Val
 995 1000 1005

Arg Glu Arg Ile Arg Gln Thr Pro Thr Ala Ala Arg Glu Arg Thr Arg
 1010 1015 1020

Gln Ala Pro Thr Ala Ala Lys Val Gly Ala Val Ser Gln Asp Leu Asp
 1025 1030 1035 1040

Lys His Gly Ala Val Ser Ser Asn Val Asn His Pro Ser Cys Ala Trp
 1045 1050 1055

Leu Glu Ala Gln Glu Glu Glu Glu Val Gly Phe Pro Glu Leu Leu Asp
 1060 1065 1070

Thr Gly Ala Asp Asp Thr Val Leu Glu Asp Ile Asn Leu Pro Gly Lys
 1075 1080 1085

Trp Lys Pro Lys Met Ile Gly Gly Ile Gly Gly Leu Ile Lys Val Lys
 1090 1095 1100

Gln Tyr Asp Gln Ile Leu Ile Glu Ile Cys Gly Lys Lys Ala Ile Gly
 1105 1110 1115 1120

Thr Val Leu Val Gly Pro Thr Pro Val Asn Ile Ile Gly Arg Asn Met
 1125 1130 1135

Leu Thr Gln Ile Gly Cys Thr Leu Asn Phe Pro Ile Ser Pro Ile Glu
 1140 1145 1150

Thr Val Pro Val Lys Leu Lys Pro Gly Met Asp Gly Pro Lys Val Lys
 1155 1160 1165

Gln Trp Pro Leu Thr Glu Glu Lys Ile Lys Ala Leu Thr Glu Ile Cys
 1170 1175 1180

Ala Asp Met Glu Lys Glu Gly Lys Ile Ser Lys Ile Gly Pro Glu Asn
 1185 1190 1195 1200

Pro Tyr Asn Thr Pro Ile Phe Ala Ile Lys Lys Lys Gln Ser Thr Lys
 1205 1210 1215

Trp Arg Lys Leu Val Asp Phe Arg Glu Leu Asn Lys Arg Thr Gln Asp
 1220 1225 1230

Phe Trp Glu Val Gln Leu Gly Ile Pro His Pro Ala Gly Leu Lys Lys
 1235 1240 1245

Lys Lys Ser Val Thr Val Leu Asp Val Gly Asp Ala Tyr Phe Ser Val
 1250 1255 1260

Pro Leu Asp Glu Ser Phe Arg Lys Tyr Thr Ala Phe Thr Ile Pro Ser
 1265 1270 1275 1280

Thr Asn Asn Glu Thr Pro Gly Val Arg Tyr Gln Tyr Asn Val Leu Pro
 1285 1290 1295

Gln Gly Trp Lys Gly Ser Pro Ile Phe Gln Ser Ser Met Thr Lys Ile
 1300 1305 1310

Leu Glu Pro Phe Arg Ser Lys Asn Pro Asp Ile Val Ile Tyr Gln Tyr
 1315 1320 1325

Met Asp Asp Leu Tyr Val Gly Ser Asp Leu Glu Ile Gly Gln His Arg
 1330 1335 1340

Thr Lys Ile Glu Glu Leu Arg Ala His Leu Leu Ser Trp Gly Phe Ile
 1345 1350 1355 1360

Thr Pro Asp Lys Lys His Gln Lys Glu Pro Pro Phe Leu Trp Met Gly
 1365 1370 1375

Tyr Glu Leu His Pro Asp Lys Trp Thr Val Gln Pro Ile Glu Leu Pro
 1380 1385 1390

Glu Lys Asp Ser Trp Thr Val Asn Asp Ile Gln Lys Leu Val Gly Lys
 1395 1400 1405

Leu Asn Trp Ala Ser Gln Ile Tyr Ala Cys Thr Pro Tyr Asp Ile Asn
 1410 1415 1420

Gln Met Leu Arg Gly Pro Gly Arg Ala Phe Val Thr Ile Pro Asn Pro
 1425 1430 1435 1440

Leu Leu Gly Leu Asp
 1445

<210> 33

<211> 4350

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Chimeric polynucleotide

<400> 33

```

ccccgccca ccatgccat cgtgcagaac gccagggcc agatgcacca ggccctgtcc 60
ccccgcaccc tgaacgcctg ggtgaagggtg atcgaggaga aggccttctc ccccgagggtg 120
atcccatgtg tctccgccct gtccgagggc gccaccccc aggacctgaa catgatgctg 180
aacatcgtgg gcggccacca ggccgccatg cagatgctga aggacaccat caacgaggag 240
gccgccgagt gggaccgcct gcaccccggtg cagccggcc ccatcccccc cgccagatg 300
cgcgagcccc gcgatccga catcgccggc accacctcca ccctgcagga gcagatcggc 360
tggatgacct ccaaccccc catccccgtg ggcgacatct acaagcgctg gatcatcctg 420
ggcctgaaca agatcgtgcg catgtactcc cccgtgtcca tcctggacat ccgccagggc 480
cccaaggagc cttccgcga ctacgtggac cgcttcttca agaccctgcg cgccgagcag 540
gccaccaggg aggtgaagaa ctggatgacc gagaccctgc tggatgcagaa cgccaacccc 600
gactgcaagt ccatcctgcg cgccctgggc cccggcgcca ccctggagga gatgatgacc 660
gcctgccagg gcgtgggcgg ccccgccac aaggcccgcg tgctgggtac cggcgcccgc 720
gcctccgtgc tgtccggcgg caagctggac gcctgggaga agatccgcct gcgccccggc 780
ggcaagaaga agtaccgcct gaagcacctg gtgtgggcct cccgcgagct ggagcgcttc 840
gccctgaacc cctccctgct ggagaccgcc gagggctgcc agcagatcat ggagcagctg 900
cagtccgccc tgaagacctc cgaggagctg aagtccctgt tcaacaccgt ggccaccctg 960
tactcgctgc accagcgcag cgacgtgaag gacaccaagg aggccctgga caagatcgag 1020
gagatccaga acaagtccaa gcagaagacc cagcaggccg ccgccgacac ccagtcctcc 1080
tccaaggtgt cccagaacta cgccctgaag caccgcgcct acgagctgga attcggcatc 1140

```

aaggtgaagc	agctgtgcaa	gctgctgcmc	ggcgccaagg	ccctgaccga	catcgtgacc	1200
ctgaccgagg	aggccgagct	ggagctggcc	gagaaccgcg	agatcctgaa	ggaccccgctg	1260
cacggcgtgt	actacgaccc	ctccaaggac	ctgatcgccg	agatccagaa	gcagggccag	1320
gaccagtgga	cctaccaa	ctaccaggag	cccttcaaga	acctgaagac	cggcaagtac	1380
gcccgaagc	gctccgccc	gaccaacgac	gtgaagcagc	tggccgaggt	ggtgcagaag	1440
gtggtgatgg	agtccatcgt	gatctggggc	aagaccccca	agttccgcct	gcccattccag	1500
aaggagacct	gggagacctg	gtggatggac	tactggcagg	ccacctggat	tcccagagtgg	1560
gagttcgtga	acacccacc	cctggtgaag	ctgtggtatc	agctggagaa	ggaccccatc	1620
gccggcgccg	agaccttcta	cgtggacggc	gccgccaacc	gcgagaccaa	gctgggcaag	1680
gccggctacg	tgaccgaccg	gggcccag	aaggtggtgt	ccctgaccga	gaccaccaac	1740
cagaagaccg	agctgcacgt	catccacctg	gcctgcagg	actccggctc	cgaggtgaac	1800
atcgtgaccg	actcccagta	cgccctgggc	atcatccagg	cccagcccga	cagatctgac	1860
cccgtggacc	ccaacctgga	gccctggaac	cacccggct	cccagcccac	caccccggc	1920
tccaagtgt	actgcaaggt	gtgctgtac	cactgccccg	tgtgcttct	gaacaagggc	1980
ctgggcatct	cctacggccg	caagaagcgc	cgccagcgcc	gcggcacc	ccagtccaac	2040
aaggaccacc	agaaccccat	ccccaagcag	cccatcccc	agaccaggg	catctccacc	2100
ggccccaagg	agtccaagaa	gaaggtggag	tccaagaccg	agaccgacc	cgaggacgcc	2160
ggccgctccg	gcaactccga	cgaggagctg	ctgaaggcca	tccgcatcat	caagatcctg	2220
taccagtcca	acccctacc	caagcccaag	ggctcccgcc	aggcccga	gaaccgccgc	2280
cgccgctggc	gcgcccggca	gcgcccagatc	gactccctgt	ccgagcgcat	cctgtccacc	2340
tgcctggggc	gccccgccc	gcccgtgccc	ctgcagctgc	ccccctgga	gctggactgc	2400
tccgaggact	gcggcacctc	cggcacccag	cagtcccagg	gcgcccagac	cggcgtgggc	2460
cgccccagg	tgtccgtgga	gtcctccgcc	gtgctgggct	ccggcaccaa	ggagggtacc	2520
gtgcgcccc	aggtgcccct	gcgccccatg	acctacaagg	ccgccttcga	cctgtccttc	2580
tttctgaagg	agaaggggcg	cctggacggc	ctgatctact	ccaagaagcg	ccaggagatc	2640
ctggacctgt	gggtgtacca	caccagggc	tacttccccg	actggcagaa	ctacaccccc	2700
ggccccggca	tccgctaccc	cctgaccttc	ggctggtgct	tcaagctggt	gcccgtggac	2760
cccagcagg	tggaggaggc	caccggcggc	gagaacaact	ccctgctgca	ccccatctgc	2820
cagcacggca	tggacgacga	ggagaaggag	acctgcgct	ggaagttcga	ctcctccctg	2880
gccctgaagc	accgcgccc	cgaactccac	cccagtcct	acaaggactg	ccccagatc	2940
accctgtggc	agcgcgccct	ggtgaccaag	atcggcggcc	agaagacgcg	tggcggcaag	3000
tggtccaagt	cctccatcgt	gggctggccc	gaggtgcgcg	agcgcatccg	ccagaccccc	3060
accgcgccc	gcgagcgcac	ccgccaggcc	cccaccgccc	ccaaggtggg	cgccgtgtcc	3120
caggacctgg	acaagcacgg	cgccgtgtcc	tccaacgtga	accacccctc	ctgcgcctgg	3180
ctggaggccc	aggaagagga	agaggtgggc	ttccccgagc	tcctggacac	cgcgccgac	3240
gacaccgtgc	tggaggacat	caacctgccc	ggcaagtgg	agcccaagat	gatcgcggc	3300
atcggcggct	tgatcaaggt	gaagcagtac	gaccagatcc	tgatcgaaat	ctgcggcaag	3360
aaggccatcg	gcaccgtgct	ggtgggcccc	accccgctga	acatcatcgg	ccgcaacatg	3420
ctgaccaga	tcggtgcac	cctgaacttc	ccatctccc	ccatcgagac	cgtgccgtg	3480
aagctgaagc	ccggcatgga	cgcccccaag	gtgaagcagt	ggccctgac	cgaggagaag	3540
atcaaggccc	tgaccgaaat	ctgcgcccag	atggagaagg	agggaagat	cagtaagatc	3600
ggccccgaga	acccctacaa	caccccatc	ttcgccatca	agaagaagca	gtccaccaag	3660
tggcgcaagc	tgggtgactt	ccgcgagctg	aacaagcgca	cccaggactt	ctgggaggtg	3720
cagctgggca	tccccaccc	cgccggcctg	aagaagaaa	agtccgtgac	cgtgctggac	3780
gtgggcgacg	cctacttctc	cgtgcccctg	gacgagtcct	tccgcaagta	caccgccttc	3840


```
accatcccct ccaccaacaa cgagaccccc ggcgtgctgct accagtacaa cgtgctgccc 3900
cagggctgga agggatcccc catcttcag tcctccatga ccaagatcct ggagcccttc 3960
cgtccaaga accccgacat cgtgatctac cagtacatgg acgacctgta cgtgggctcc 4020
gacctggaga tcggccagca ccgcaccaag atcgaggagc tgcgcgccca cctgctgtcc 4080
tggggcttca tcacccccga caagaagcac cagaaggagc cccccttcct gtggatgggc 4140
tacgagctgc accccgacaa gtggaccgtg cagcccatcg agctgcccgga gaaggactcc 4200
tggaccgtga acgacatcca gaagctggtg ggcaagctga actgggcctc ccaaactctac 4260
gcctgcaccc cctacgacat caaccagatg ctgcgcggcc ccggccgcgc cttcgtgacc 4320
atccccaacc ccctgctggg cctggactag 4350
```